

**A PARTIAL DESCRIPTION OF THE PARAMETER SPACE
OF RATIONAL MAPS OF DEGREE TWO: PART 2**

Mary Rees

Department of Pure Mathematics, University of Liverpool,
P.O. Box 147, Liverpool L69 3BX, U. K.

Address until 30th June 1991:
Institute of Mathematical Sciences, SUNY at Stony Brook,
Stony Brook NY 11794-3660, U.S.A.

Abstract

This continues the investigation of a combinatorial model for the variation of dynamics in the family of rational maps of degree two, by concentrating on those varieties in which one critical point is periodic. We prove some general results about nonrational critically finite degree two branched coverings, and finally identify the boundary of the rational maps in the combinatorial model, thus completing the proofs of results announced in Part 1.

Introduction.

This is the second in a series of papers aimed at understanding the variation in dynamical behaviour within the family of rational maps of degree two. This introduction will be brief, as Chapter 1 of [R2] was intended as an introduction to this paper also, and includes statements of the results of this paper.

As in [R2], we concentrate on maps which are *critically finite*. Recall that an orientation-preserving branched covering $g : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ is called *critically finite* if $\#(X(g)) < \infty$, where

$$X(g) = \{g^n(c) : c \text{ critical, } n > 0\},$$

and that these play an important rôle in the understanding of rational maps. A branched covering of $\bar{\mathbf{C}}$ of degree two necessarily has two critical points c_1, c_2 . More generally, we consider degree two branched coverings for which one critical point is periodic. Likewise, these play an important rôle. Douady and Hubbard have studied the *Mandelbrot set* for degree two polynomials (for which the critical point at ∞ is, of course, fixed). They provided a combinatorial model for the Mandelbrot set ([D-H1], [D-H2]) which was reinterpreted by Thurston [T]. The work described in this paper, and its predecessor and successors, is a generalisation of the combinatorial model of the Mandelbrot set to maps with a critical point which is no longer fixed, but still periodic. It seems likely that this model can be completed to include maps for which both critical points have infinite forward orbits.

Recall that two critically finite branched coverings f, g are *equivalent*, written $f \simeq g$, if there exists an orientation-preserving homeomorphism $\varphi : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ and a path g_t ($t \in [0, 1]$) through critically finite branched coverings such that $X(g_t) = X(g)$ for all $t \in [0, 1]$, $\varphi \circ f \circ \varphi^{-1} = g_0$, $g = g_1$. Thurston [T] gave a necessary and sufficient condition for a critically finite branched covering to be equivalent to a rational map. Almost always - and always in degree two - there is at most one rational map in an equivalence class, and the rational map can be determined up to topological conjugacy (and its Julia set, up to homeomorphism) by any map in the equivalence class. More detail is given on this in [R2].

We are interested in understanding rational maps, but, paradoxically, the subject of this paper is, very largely, critically finite branched coverings which are not equivalent to rational maps. The reason is basically the following. Roughly speaking, it is easier to describe dynamically the family of all critically finite degree two branched coverings with, say, one critical point of period m than simply the family of all such rational maps. (We remark that there is not so much difference between the two families if $m = 1$.) It is also easier to analyse nonrational branched coverings (up to equivalence) rather than rational maps, and then, of course, the rational ones are what remain when the nonrational ones are removed.

The chapters of this paper give increasingly precise results about increasingly specified branched coverings. The last two chapters concern lamination

maps, which are, for the present, our preferred models for rational maps. Their dynamics are easily computable, and a lamination map determines an equivalent rational map up to topological conjugacy. This is the substance of the Lamination Map Conjugacy Theorem in [R2]. The chapters, and their rough content, are as follows.

Chapter 1: Homotopy Information about Branched Coverings of Degree Two.

Thurston's theorem giving a necessary and sufficient condition for equivalence of a critically finite branched covering f to a rational map ([T], [D-H3]) is in terms of the action of f^{-1} on isotopy classes of loops in $\bar{\mathbb{C}} \setminus X(f)$. This condition, and a couple of refinements in degree two, which are essentially isotopy-invariance conditions, are stated in the introduction of [R2]. This chapter examines more closely the structure of f when f is *not* equivalent to a rational map. In particular, a *critical branched covering* is associated to f (1.9), essentially a restriction of an iterate of f to a subsurface. It may be degree one or two, and is in some sense irreducible. The main result of the chapter is the Decomposition Proposition of 1.12.

Chapter 2: The Fixed Holed-Sphere Theorem.

The isotopy-invariance of the first chapter is translated into actual invariance, in the Fixed Holed-Sphere Theorem. Underlying this is an application of the common idea that a map with homotopy type satisfying certain “expanding” or “hyperbolic” properties is determined up to a useful semiconjugacy by its homotopy type.

Chapter 3: 1-dimensional Topological Dynamics.

No particular originality is claimed for this. The 1-sided Neighbourhood Theorem is similar (though not identical, so far as I know) to many in the literature, and shows, roughly, that under certain mild conditions, a graph preserved by a map is a union of unstable manifolds.

Chapter 4: The Nonrational Lamination Map Theorem.

Now we specify to lamination maps, and show that an invariant lamination with a critically finite lamination map with a nondegenerate Levy cycle (roughly speaking, not equivalent to a rational map) contains periodic leaves - with a specified bound on the period - with certain additional properties. Finally, this result is used in

Chapter 5: Proof of the Admissible Boundary Theorem.

Lamination maps (at least, those which have been defined at any point in this work) have 0 as a periodic critical point, and lie in countably many families, depending on the definition of the map on a set containing the orbit of 0. The lamination maps in each family are identified with leaves or complementary components - called gaps - of a lamination on a disc, called a parameter lamination and denoted by $\mathcal{L}_r$, where r runs through the odd denominator rationals in $(0, 1)$. For details of this structure, see Chapter 1 of [R2]. This chapter contains the proof of two results stated in the introduction of [R2] - the Admissibil-

ity Proposition and the Admissible Boundary Theorem. Identifying lamination maps with the corresponding leaves and gaps in the laminated disc, these two theorems together identify an open set of lamination maps (and its boundary). In this set, all critically finite branched coverings have at most only degenerate Levy cycles, which virtually means they are all equivalent to rational maps, and the set contains all critically finite rational maps, up to equivalence.

Chapter 1. Homotopy information about Branched Coverings of Degree Two.

1.1.

The information in this chapter is not very deep, and most of it is known anyway: it is repeated for convenience. Let $f : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ be a critically finite degree two orientation-preserving branched covering. Then exactly one of the following occurs, as can be extracted from [T], and is explained in 1.2 - 1.3 below.

Up to equivalence, f is a rational map.

Up to equivalence, f is $[x] \mapsto [Ax + b] : T^2 / \sim \rightarrow T^2 / \sim$, where T^2 is the two-torus, $\sim$ is the relation $x \sim -x$, $[x]$ is the equivalence class of x , $b \sim -b$ and A is an automorphism of T^2 with determinant 2 and eigenvalues λ_s, λ_u , $|\lambda_s| < 1 < 2 < |\lambda_u|$.

There is a Levy cycle for f . (This term is recalled below.)

The third possibility is in some sense a decomposable case. We shall define a *complete Levy set* for f , consisting of a union of disjoint Levy cycles and some preimages of these. Given f and a complete Levy set Γ , we shall see that components of $\bar{\mathbf{C}} \setminus \cup \Gamma$ are eventually periodic in a natural sense, and hence have maps associated to them, which are always either homeomorphisms or degree two critically finite branched coverings. In particular, such a map is associated to the component of $\bar{\mathbf{C}} \setminus \cup \Gamma$ containing both critical points, in the case when the forward orbits of both critical points of f contain periodic critical points, and this map is of one of the first two indecomposable types if it is degree two. We shall summarise most of the information in the Decomposition Proposition (1.12). This information will be crucial in the semiconjugacy results for critically finite degree two branched coverings, in Chapter 2. It is clear that a more complete description of those critically finite degree two branched coverings with Levy cycles is possible, partly in terms of a natural tree obtained by treating Levy loops as vertices, and partly in terms of the maps associated with components of $\bar{\mathbf{C}} \setminus \cup \Gamma$. I suppose the description would be reminiscent of the Douady-Hubbard-Thurston description of critically finite quadratic polynomials, with some additional identifications. In fact, Shishikura [S] has some results in a more general framework: he has constructed a tree (with an expanding metric) using Thurston obstruction loops as vertices.

1.2. The cases when a Thurston obstruction does not exist.

The Thurston obstruction is a necessary and sufficient condition for nonequivalence to a rational map, when a certain orbifold Q_g , associated to an orientation-preserving critically finite branched covering g , is *hyperbolic* ([T] Ch. 16). Without going into too much detail about what this means, we identify those g of

degree two which are not equivalent to rational maps, and for which Q_g is not hyperbolic. If Q_g is not hyperbolic, then $\#(X(g)) \leq 4$ ([T] Ch. 16).

If g and h are critically finite orientation-preserving branched coverings and $\#(X(g)) \leq 3$, then g and h are equivalent if and only if there is an isomorphism between $(g^{-1}X(g), g)$ and $(h^{-1}X(h), h)$ which maps critical points to critical points of the same multiplicity. This follows from the fact that any homeomorphism of the sphere which fixes three points is isotopic to the identity, via an isotopy fixing those three points. From this, it is easy to see that if $\#(X(g)) \leq 3$, g is equivalent to a rational map.

If $\#(X(g)) = 4$ and Q_g is not a hyperbolic orbifold, then there is only one possibility for the universal orbifold cover $\tilde{Q}_g$ of Q_g and for the covering map $\pi : \tilde{Q}_g \rightarrow Q_g$: $\tilde{Q}_g = \mathbf{C}$, Q_g is the quotient of $\mathbf{C}$ by the relation $z \sim \pm z + m + ni$ ($m, n \in \mathbf{Z}$) and π is the quotient map. Since g^{-1} lifts to Q_g and induces a homomorphism of a subgroup of the covering group onto the covering group (as always: see [T] 16.6) and the covering group in this case is of the form $\mathbf{Z}^2 \times_s \mathbf{Z}_2$, we see that g^{-1} lifts injectively in this case, so that g also lifts - to $\tilde{g} : \mathbf{C} \rightarrow \mathbf{C}$, preserving equivalence classes in general, and those of critical orbits in particular. Hence, g is equivalent to $[z] \mapsto [Az + \beta]$, for some real-linear map A and $\beta \in (\mathbf{Z}/2)^2$. In addition, $\det(A) = \text{degree}(g)$. Let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

If $(a + d)^2 < 4(ad - bc)$, A is conjugate to a skew-symmetric matrix, and g is equivalent to a rational map. Now let $ad - bc = 2 = \text{degree}(g)$. If $(a + d)^2 = 9$, then the existence of an eigenvector of A with eigenvalue ± 1 implies the existence of a simple loop γ in $\bar{\mathbf{C}} \setminus X(g)$ such that a component γ' of $g^{-1}(\gamma)$ is isotopic to γ in $\bar{\mathbf{C}} \setminus X(g)$, and $g|_{\gamma'}$ is a homeomorphism. If $(a + d)^2 > 9$, then A has eigenvalues λ_s, λ_u with $|\lambda_s| < 1 < 2 < |\lambda_u|$.

1.3. Levy Cycles.

Let $g : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ be a critically finite degree two orientation-preserving branched covering. Then we recall that a set $\{\gamma_1 \dots \gamma_r\}$ of simple nontrivial disjoint loops in $\bar{\mathbf{C}} \setminus X(g)$ is a *Levy cycle for g* if there exists γ'_i in $g^{-1}(\gamma_{i+1})$ (writing $\gamma_{r+1} = \gamma_1$) such that γ_i and γ'_i are isotopic in $\bar{\mathbf{C}} \setminus X(g)$ and $g|_{\gamma'_i}$ is a homeomorphism. We recall that Thurston's theorem for a necessary and sufficient condition for *non-equivalence* to a rational map [T] was given three equivalent formulations in the case of degree two in 1.6 of [R2] (where further references are given). The middle condition was the existence of a Levy cycle. In [R2] the hypothesis of Thurston's theorem was slightly simplified. In fact, the theorem holds whenever the associated orbifold is hyperbolic. Non-hyperbolic orbifolds have been investigated in 1.2, and, putting these facts together, we

see that, if g is an orientation-preserving degree two critically finite branched covering, exactly one of the three properties stated in 1.1 does indeed hold.

So now we wish to obtain more information about g , where g is a degree two critically finite orientation-preserving rational map with a Levy cycle. We start with two lemmas, which are, in fact, the basis of a proof of existence of a Levy cycle in the first place ([TL], for example). The proofs are included for convenience, since they are quite short.

1.4. Lemma. *Let $g : \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$ be a critically finite degree two orientation-preserving branched covering. Let Γ be a (finite) set of simple disjoint isotopically distinct loops in $\bar{\mathcal{C}} \setminus X(g)$ with $\Gamma \subset g^{-1}\Gamma$ up to isotopy. Let $\gamma \in \Gamma$ separate critical values. Then γ bounds a disc D such that all components of $g^{-n}D$ are discs, for all $n \geq 0$.*

Proof:

Let Γ_1 be the set of loops γ in Γ such that γ bounds a disc D for which all components of $g^{-n}D$ are discs for all $n \geq 0$. Then $\Gamma \cap g^{-1}\Gamma \subset \Gamma_1$. (We mean “up to isotopy in $\bar{\mathcal{C}} \setminus X(g)$ ”, but we omit saying this, for the rest of this proof.) Let $\Gamma_2 = \Gamma \setminus \Gamma_1$. Then $\Gamma_2 \subset g^{-1}\Gamma_2$. We claim that no loop in Γ_2 separates critical values. Otherwise, let D be a disc containing exactly one critical value, bounded by a loop γ from Γ_2 , and containing no loop from Γ_2 in its interior. Then we can prove by induction on n that all components of $g^{-n}D$ are discs, and none contains a loop from Γ_2 in its interior. For assume this is true for n , and D_1 is a disc component of $g^{-n}D$. Then D_1 cannot contain both critical values, since it does not contain γ in its interior. So components of $g^{-1}D_1$ are all discs, and do not contain elements of $g^{-1}\Gamma_2 \supset \Gamma_2$ in their interiors. The inductive step is completed. So $\gamma \in \Gamma_1$, giving a contradiction, and the lemma is proved.

1.5. Degenerate Levy Cycles.

A Levy cycle $\Gamma = \{\gamma_1 \dots \gamma_r\}$ for g is *degenerate* if each γ_i, γ'_i (as in 1.3) bounds a disc D_i, D'_i , with D'_i isotopic to D_{i+1} via an isotopy preserving X , and $g : D'_i \rightarrow D_i$ is a homeomorphism. Thus no disc D_i contains a critical value of g . So if, for instance, the forward orbits of both critical points of g contain periodic critical points, any Levy cycle for g must be *nondegenerate*. The following lemma is then immediate.

Lemma.

Let g be a critically finite degree two orientation-preserving branched covering, such that any Levy cycle is nondegenerate. Let γ be a loop in $\bar{\mathcal{C}} \setminus X(g)$ bounding a disc D such that all components of $g^{-n}D$ are discs for all $n \geq 0$. Then D cannot contain any loop of a Levy cycle.

1.6. Complete Levy sets.

Let $f : \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$ be a critically finite degree two orientation-preserving branched covering, and let Γ be a finite set of disjoint simple nontrivial loops in $\bar{\mathcal{C}} \setminus X(f)$. Then Γ is a *complete Levy set* (for f) if the conditions below hold, writing $X = X(f)$.

The Levy Condition. $\Gamma \subset f^{-1}\Gamma$, and for $\gamma \in f^{-1}\Gamma$, $\gamma \in \Gamma$ if and only if γ does not bound a disc D such that all components of $f^{-n}D$ are discs, for all n .

Thus, by Lemma 1.4, no loop in Γ separates critical values, and f is degree one restricted to any component of $f^{-1}\Gamma$. There are Levy cycles in Γ , but Γ may not be a union of Levy cycles.

The Gap Invariance Condition. If G is a component of $\bar{\mathcal{C}} \setminus \cup \Gamma$ such that $G \setminus X$ is not an annulus, then there is a unique component G' of $\bar{\mathcal{C}} \setminus X$ - which we term f_*G - and a unique component G'' of $f^{-1}G'$ such that $G'' \subset G$ and $\partial G \subset \partial G''$.

Definition. G is a *gap* for Γ if G is a component of $\bar{\mathcal{C}} \setminus \cup \Gamma$ such that $G \setminus X$ is not an annulus.

The Gap Restrictions. No annulus gap of $\bar{\mathcal{C}} \setminus (X \cup (\cup \Gamma))$ is mapped onto itself by f^n (any $n > 0$). If G_1, G_2 are distinct with $\bar{G}_1 \cap \bar{G}_2 \neq \emptyset$, then at least one G_i is not mapped homeomorphically by all f^n . Any Levy cycle disjoint from Γ can be isotoped in $\bar{\mathcal{C}} \setminus X$ either into Γ itself or into a periodic cycle of gaps on which f acts homeomorphically.

1.7. Lemma. *Let g have at least one nondegenerate Levy cycle, but no degenerate ones. Then there exists a complete Levy set Γ for some f , where f is homotopic to g via a homotopy $\{f_t\}$ through critically finite branched coverings with $X(f_t) = X(g)$ for all t .*

Proof: Let Γ_0 be a maximal union of disjoint Levy cycles for g . Let $k > 0$ be such that, if $\gamma \in g^{-(k+1)}\Gamma_0$, and γ does not bound a disc D such that all components of $g^{-n}D$ are discs for all $n > 0$, then $\gamma \in g^{-k}\Gamma_0$ up to isotopy. Let Γ_1 be a set of two copies, up to isotopy, of all such loops in $g^{-(k+1)}\Gamma_0$ (or, equivalently, in $g^{-k}\Gamma_0$). Then g, Γ_1 satisfy the Levy Condition except that $\Gamma_1 \subset g^{-1}\Gamma_1$ up to isotopy only. Note that Γ_1 contains up to isotopy all loops in $g^{-1}\Gamma_1$ which separate two elements of Γ_1 . Hence G'', G' exist as in the Gap Condition, up to isotopy. Then f can be chosen equivalent to g (via a homotopy preserving X) so that the Levy and Gap Conditions are satisfied by f, Γ_1 . If G_1, G_2 are distinct gaps for Γ_1 such that f^n always maps both of them homeomorphically, then if one of them is periodic they both are, and the period of both is $\leq$ the period of the common boundary component. So obtain Γ from Γ_1 by removing all such common boundary components and all their preimages under all f^n , and also removing redundant loops in annuli (possibly removing some annuli gaps altogether) so that Γ satisfies the Gap Restrictions.

The last Gap Restriction is satisfied because Γ_0 was a maximal union of Levy cycles.

1.8. Critical gaps for Complete Levy Sets.

From now on, until the end of this chapter, let $f : \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$ be a critically finite degree two orientation-preserving branched covering with no degenerate Levy cycle, and let Γ be a complete Levy set for f . Then by Lemma 1.4, and the Levy Condition, both critical values of f are contained in the same gap for Γ . Hence, both critical points are contained in the same gap, which we call the *critical gap*, usually denoted by G_0 . The following lemma holds, under the hypotheses of this section. It, too, can be found essentially in [TL].

Lemma. *The disc gaps for Γ are $f_*^i G_0$, with $1 \leq i \leq r$ for some $r \geq 2$.*

Proof: There is a gap G which is a disc. Then $G \cap X(f) \neq \emptyset$. So $f^n c \in G$ for some least $n \geq 1$ and c critical. If $n > 1$ then both components of $f^{-1}G$ are discs. Then if we write $G_n = G$, we can find G_i ($1 \leq i \leq n$) with $f_* G_i = G_{i+1}$ for $i < n$ and $f^i c \in G_i$. Then G_i is a disc for each i , and G_1 is the gap containing the critical values, as required. We must have $r \geq 2$, because $\bar{\mathcal{C}}$ has Euler characteristic 2, so at least two components of $\bar{\mathcal{C}} \setminus (\cup \Gamma)$ are discs.

1.9. Branched coverings associated to gaps.

Let G be a gap for Γ which is of period n under f_* . (All gaps must be eventually periodic under f_* , since there are only finitely many gaps.) Let G_1 be the component of $f^{-n}G$ with $G_1 \subset G$, $\partial G \subset \partial G_1$. Then $G_1 = SG$ for an inverse branch S of the branched covering f^n . Inductively we can define $G_{k+1} = SG_k$. Then for some least $k \geq 0$, all components of $\partial G_{k+1} \setminus \partial G_k$ are trivial in $\bar{\mathcal{C}} \setminus X(f)$.

Now f^n maps components of ∂G_{k+1} to components of ∂G_k , and maps components of ∂G to components of ∂G . Also, $f^n|_{G_{k+1}}$ is either a homeomorphism or a degree two branched covering, depending on whether or not the critical gap is in the forward orbit of G . Then we define $h = h_G$ by

- a) $h = f^n$ on G_{k+1} ,
- b) $h(D) = D'$ if D, D' are components of $\bar{\mathcal{C}} \setminus G_{k+1}, \bar{\mathcal{C}} \setminus G_k$ with $h(\partial D) = \partial D'$,
- c) each component of $\bar{\mathcal{C}} \setminus G_{k+1}$ contains at most one point of $h^{-1}X(h)$.

Then if G is in the forward orbit of the critical gap, h is the *critical branched covering* of f . It is uniquely defined up to equivalence (given Γ), and h has no Levy cycle, since Γ is a complete Levy set. (See the Gap Restrictions of 1.6.) If the critical gap is not in the forward orbit of G , then $k = 1$ and $f^n = h : G \rightarrow G$ is a homeomorphism.

1.10. Lemma. *Let the critical gap G_0 of Γ be periodic and the critical branched covering h be not equivalent to a rational map. Then the period of G_0 is less than the period of any periodic critical point of f .*

Proof: Let n be the period of G_0 . Let G_1 be the component of $f^{-2n}G_0$ with $G_1 \subset G_0$, $\partial G_0 \subset \partial G_1$. Now by 1.2, h has no fixed critical points (in fact, no periodic critical points). Then for any critical point c , there is a disc component D of $G_0 \setminus G_1$ containing c such that $f^n D$ is another component of $G_0 \setminus G_1$ and $f^i D \cap G_0 = \emptyset$ for $0 < i < n$. So if c is periodic, c must have period $> n$.

1.11. The Fixed Set P .

By the Lefschetz Fixed Point Theorem, some loop or gap P of Γ is fixed by f . (This is the third equivalent condition of 1.6 of [R2] for nonequivalence to a rational map. This proof is due to Shishikura.) If P is a loop, it must have orientation reversed, because otherwise one of the discs it bounds is fixed, contradicting the Levy Condition. Neither can P be the critical gap, because if it is, any periodic boundary component is a degenerate Levy cycle for f . So if P is a gap, $f : P \rightarrow P$ is a homeomorphism, and boundary components are cyclically permuted, again because there are no degenerate Levy cycles.

1.12.

Now we can summarise the main homotopy-type information we shall use.

Decomposition Proposition.

Let $g : \bar{C} \rightarrow \bar{C}$ be an orientation-preserving critically finite degree two branched covering which is not equivalent to a rational map, and with no degenerate Levy cycle. Let f be an equivalent branched covering (via a homotopy which is constant on $X(g) = X(f)$) with complete Levy set Γ , critical gap G_0 and fixed loop or gap P . Let ℓ be the number of boundary components of P , or $\ell = 2$ if P is a loop. Then one of the following holds.

Rational Critical Branched Covering. *The critical gap G_0 is periodic under f_* , and the critical branched covering is equivalent to a rational map.*

Periodic Separating Set. *A gap Q is mapped homeomorphically onto itself by f^ℓ , Q separates P and f_*G_0 , and $\bar{Q} \cap \bar{P} = \bar{Q} \cap \overline{f_*G_0} = \emptyset$.*

Adjacent Periodic Set. *The gap f_*G_0 is periodic, and adjacent to some periodic gap or loop of lesser period than itself.*

1.13. Outline of proof of the Decomposition Proposition.

Now we fix some notation. Let Γ be a complete Levy set for f , and let P , ℓ be as in the statement of the proposition. Let R be the set bounded by P and f_*G_0 - which is either a closed annulus or a loop of Γ . Clearly, if R is a loop of Γ , G_0 has period ℓ , and the converse is also true, because no annulus component of $\bar{\mathbf{C}} \setminus \cup \Gamma$ can be mapped homeomorphically to itself by f^ℓ , by the definition of complete Levy set. Let $\gamma = \partial R \cap \partial P$. Let T be the local inverse of f^ℓ defined on R such that T preserves γ . Let S be the local inverse of f^ℓ defined on R which is $\neq T$ and such that SR is in the same component of $\bar{\mathbf{C}} \setminus P$ as f_*G_0 and R . Now we prove the following lemma, under the assumptions that R is a closed annulus and G_0 does not have period ℓ . It clearly completes the proof of the Decomposition Proposition.

1.14. Lemma. *$TR \subset R$, $SR \cap R \neq \emptyset$, and there is a set Q in $SR \cap R$ which is either a gap or loop of Γ , such that Q has period ℓ under f and is mapped homeomorphically by f^ℓ , and Q separates f_*G_0 from TR .*

We split the proof of this lemma itself into a number of stages.

1.15. Lemma. *There are least integers N , $N' \geq 0$ with*

$$f_*^{N\ell+1}G_0 \cap SR = \emptyset, \quad f_*^{N'\ell+1}G_0 \cap TR = \emptyset.$$

Proof: The property derived from the definition of complete Levy set that we use repeatedly is:

- (1) for all $i \geq 0$, $f_*^iG_0$ is the connected union of a component of $f^{-1}f_*^{i+1}G_0$ and some (possibly none) discs which are disjoint from $\cup \Gamma$.

If N does not exist (for example) then $f_*^iG_0$ is in the component of $\bar{\mathbf{C}} \setminus P$ bounded by $f^{i-1}\gamma$ for all i . (If N does exist, this is true for $i \leq N\ell$.) If G_0 is periodic, this implies $G_0 = f_*^{p\ell}G_0$ for some $p \geq 2$. But $G_0 \cap f^{-1}R = \emptyset$, so then $f_*^{(p-1)\ell+1}G_0 \cap SR = \emptyset$, and N exists. If G_0 is not periodic, there are integers $i, j \geq 1$ such that $f_*^{i\ell+1}G_0 = f_*^{j\ell+1}G_0$ but $f_*^{i\ell}G_0, f_*^{j\ell}G_0$ are distinct and separated by G_0 . Then $f_*^{i\ell}G_0, f_*^{j\ell}G_0$ cannot intersect the same component of $f^{-1}R$. Then $f_*^{(i-1)\ell+1}G_0$ or $f_*^{(j-1)\ell+1}G_0$ is disjoint from SR , and, again, N exists. The proof of the existence of N' is exactly similar.

Now $f_*^iG_0 \neq f_*^{N\ell+1}G_0$ for $1 < i < N\ell + 1$, so $f_*^iG_0$ ($1 < i \leq N\ell + 1$) are all distinct. Hence they are all disjoint, and disjoint from P for $0 \leq i < N\ell + 1$. A similar statement holds with N replaced by N' .

1.16. Lemma. *$N' = 0$, so that $TR \subset R$ and $SR \cap R \neq \emptyset$, and $S\partial f_*G_0$ separates f_*G_0 from P .*

Proof: It follows from 1.15 that

- (1) $f_*^{i\ell+1}G_0$ is the union of $Tf_*^{(i+1)\ell+1}G_0$ and some discs disjoint from $X(f)$, for $0 \leq i \leq N' - 2$.

Let $R'_0 = R$. Inductively, define $R'_{i+1} = TR'_i \setminus f_*G_0$. Then $R'_i \neq \phi$ for all i by induction, since $\gamma \subset TR'_i$, and $R'_{i+1} \subset R'_i$. But by 1.15, using (1),

$$TR'_j \cap f_*^{i\ell+1}G_0 \neq \phi, \text{ but } TR'_j \cap f_*^{(N'-j)\ell+1}G_0 = \phi$$

for $0 \leq j \leq N'$, and $0 \leq i < N' - j$. In particular, $TR'_i \cap f_*G_0 \neq \phi$ for $0 \leq i < N'$, but $TR'_{N'} \cap f_*G_0 = \phi$, so $R'_{N'+1} = TR'_{N'}$, and

$$\partial R'_j = \gamma \cup \bigcup_{i=0}^j T^i \partial f_*G_0 \text{ for } j \leq N',$$

$$\partial R'_{N'+1} = \gamma \cup \bigcup_{i=1}^{N'+1} T^i \partial f_*G_0.$$

See Diagram 1. The shaded region denotes $TR_{N'}$ (drawn in the case $N' = 1$, which does not actually happen).

Diagram 1

Then $T^{N'+1}\partial f_*G_0$ must bound a disc disjoint from $R'_{N'+1} = TR'_{N'}$ containing ∂f_*G_0 . Then all boundary components of $T^k R'_{N'}$ are nontrivial in $\bar{\mathbf{C}} \setminus X$ for $k \geq 0$, $T^{k+1}R'_{N'} \subset T^k R'_{N'}$ and for some $p \geq 0$, $T^{p+1}R'_{N'}$ and $T^p R'_{N'}$ are

isotopic in $\bar{\mathbf{C}} \setminus X(g)$. If $N' > 0$ then $T^p R'_{N'}$ has $N' + 2 \geq 3$ boundary components, all in the full orbit of ∂R . Since Γ is a complete Levy set, $\partial T^p R'_{N'} \subset \Gamma$, and there is a gap G for Γ isotopic to $T^p R'_{N'}$. But then $\bar{G} \cap \bar{P} \neq \phi$ with $f^n G \neq G_0$ for any $n \geq 0$, which is a contradiction to the definition of a complete Levy set. So $N' = 0$. In particular, $TR \subset R$. Then TR and SR are separated by a component of $f^{-\ell} f_* G_0$ which is disjoint from $f_* G_0$. So $SR \cap R \neq \phi$, and $S\partial(f_* G_0)$ separates $f_* G_0$ from P .

1.17. Proof of Lemma 1.14.

Let $R = R_0$, and let R_1 be the annulus bounded by $S\partial(f_* G_0)$ and $\partial f_* G_0$, so that $R_1 \subset R = \text{domain}(S)$, and $R_1 = R \cap SR$. Then define

$$R_{i+1} = SR_i \setminus f_* G_0.$$

As in 1.16, we see that

$$SR_j \cap f_*^{i\ell+1} G_0 \neq \phi, \text{ but } SR_j \cap f_*^{(N-j)\ell+1} G_0 = \phi$$

for $0 \leq j \leq N$ and $0 \leq i < N - j$. Hence, $R_j \neq \phi$ for $0 \leq j \leq N + 1$, $R_{N+1} = SR_N$. Then if $N \geq 1$,

$$\partial R_j = \cup_{i=0}^j S^i \partial G_0 \text{ for } j \leq N,$$

$$\partial R_{N+1} = \cup_{i=1}^{N+1} S^i \partial G_0.$$

If $N = 0$, $\partial R_1 = S\partial R$. Proceeding as in the case of TR , we obtain Q . If $N \geq 2$, Q is isotopic to $S^p R_N$ for some $p \geq 0$ and is a gap for Γ of period ℓ with boundary components cyclically permuted. If $N = 0$ or 1 , Q is isotopic to $\partial S^p R_N$ for $p \geq 0$, Q is part of a Levy cycle of period ℓ and has oriented period 2ℓ , up to isotopy. So Q has the required properties.

1.18 Note. Tan Lei's proof of the existence of a minimal Levy cycle [TL] (that is, the existence of P in the current language) was obtained by taking the least integer $n > 1$ with $f_*^n G_0 \cap P' = \phi$, where P' is the annulus separating $f_* G_0$ and $f^{-1} f_* G_0$, and then taking successive inverse images under f^n of $P' \setminus \cup_{i < n} f_*^i G_0$. The proof above was obtained by generalising this idea.

1.19. Examples.

It might be as well to give some examples to show that the various alternatives of the Decomposition proposition are all quite common. We use matings to obtain the examples, and refer to 1.19 of [R2] for the notations. So let p, q be odd denominator rationals. Then $f = s_p \amalg s_q : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ is a type IV branched covering which preserves a lamination $L_p \cup L_q^{-1}$. If f is not a rational map, then some leaves of $L_p \cup L_q^{-1}$ join up to form closed loops.

Example 1. If $p = q = 1/3$, the two leaves joining the points $1/3, 2/3$ join to give a loop which is, in itself, a Levy cycle, with the critical gap adjacent to it. We can easily identify the critical branched covering by examining $(s_p \amalg s_q)^2$ restricted to the union of the gaps containing $0, \infty$, and we see that the critical branched covering is equivalent to $z \mapsto z^2$. This gives the Rational Critical Branched Covering alternative of the Decomposition Proposition.

Example 2. If p and $1 - q$ are separated from 0 by a finite-sided gap of QML - such as the triangle with vertices at $9/56, 11/56, 15/56$ - but p and $1 - q$ are not in the same component of the finite-sided gap complement, then the critical branched covering is a homeomorphism.

Example 3. Let p, q be on minor leaves which are separated from 0 by the leaf with endpoints $\frac{17}{48}, \frac{19}{48}$. See Diagram 2.

Diagram 2

As in example 1, we can identify the critical branched covering by examining $(s_p \amalg s_q)^2$ restricted to the union of the gaps containing $0, \infty$, and in this case, the critical points c_1, c_2 of the critical branched covering h satisfy $h(c_1) \neq h(c_2)$, but $h^2(c_1) = h^2(c_2)$, and $h^3(c_i) \neq h^2(c_i)$ but $h^4(c_i) = h^3(c_i)$. Then h is actually equivalent to a rational map. But we can modify this example as follows. First, perturb $s_p \amalg s_q$ to g having a complete Levy cycle, so that the critical gap G contains a four-holed sphere component G' of $g^{-2}G$ not intersecting $X(g)$. Then we can change g to another branched covering g_1 with $X(g) = X(g_1)$ by changing the definition only on $G' \cup gG'$. Then we can choose g_1 so that the new

critical branched covering is equivalent to $[x] \mapsto [Ax] : T^2 / \sim \rightarrow T^2 / \sim$, where A is any integral matrix with determinant 2 and all matrix entries odd, to ensure that the critical orbits have the same pattern as for h .

Chapter 2. The Fixed Holed-Sphere Theorem.

2.1.

In this chapter we prove a semiconjugacy result. We recall that there are many instances in dynamical systems of homotopy information (usually containing some sort of “expanding” or “hyperbolic” criterion) for a map g yielding semiconjugacy information, that is, the existence of a nontrivial map g_1 and continuous φ such that $\varphi \circ g = g_1 \circ \varphi$. The reference which comes to mind, and which is actually relevant here, is [Fr]. There it is proved (for example) that if T^2 is the two-torus, and $g : T^2 \rightarrow T^2$ is continuous and homotopic to a linear automorphism g_1 of T^2 with both eigenvalues off the unit circle, then a φ homotopic to the identity exists. The same result holds - and can be proved more simply - if g_1 is, instead, an expanding linear endomorphism. However, somewhat surprisingly, the result does not hold if g_1 is simply an endomorphism, even if both eigenvalues are off the unit circle. We can see this as follows. Let g_1 be such that the associated matrix A is in $GL(2, \mathbf{Z})$ and has determinant 2 and eigenvalues λ_u, λ_s with $|\lambda_s| < 1 < 2 < |\lambda_u|$, with corresponding eigenvectors v_u, v_s . Identify the two-torus with $\mathbf{R}^2/\mathbf{Z}^2$. We are now going to construct a covering map $g : T^2 \rightarrow T^2$ by constructing the well-defined lift, $\bar{g}$, of the multivalued function g^{-1} to $\mathbf{R}^2$ (so that notation agrees with what we shall do later). In fact, we can take $\bar{g}$ to be an arbitrarily small C^0 perturbation of A^{-1} . We choose $\bar{g}$ so that $\bar{g}(0,0) = (0,0)$, and for some $(m,n) \in \mathbf{Z}^2$, $(m,n) \neq (0,0)$, the set

$$\{(x,y) : \lim_{k \rightarrow \infty} \bar{g}^k(x,y) = (0,0)\}$$

contains (m,n) . Note that the set

$$\{(x,y) : \lim_{k \rightarrow \infty} A^{-k}(x,y) = (0,0)\}$$

is simply $\text{sp}(v_u)$, hence is not rationally defined and does not contain (m,n) .

Then if Franks' theorem held for g , there would exist $\Psi : \mathbf{R}^2 \rightarrow \mathbf{R}^2$ satisfying

- a) $\Psi((x,y) + (p,q)) = \Psi(x,y) + (p,q)$ for all $(x,y) \in \mathbf{R}^2$ and all $(p,q) \in \mathbf{Z}^2$,
- b) $\Psi \circ \bar{g} = A^{-1} \circ \Psi$.

Note that b) yields $\Psi(0,0) = (0,0)$. Then $\Psi(\bar{g}^k(m,n)) = A^{-k}(m,n) \rightarrow 0$ as $k \rightarrow \infty$, giving a contradiction to the existence of Ψ . Despite the fact that Franks' theorem does not hold in such circumstances, the technique of the proof can still be used to obtain weaker results. We shall do this in the case we are interested in, namely, branched coverings of $\bar{\mathbf{C}}$. Recall that, in Chapter 1, torus endomorphisms came up in the description of branched coverings of $\bar{\mathbf{C}}$. This is the reason for the rather involved statement of the Fixed Holed-Sphere Theorem below. Note that the theorem has two alternative conclusions. Awkward torus maps (such as the one above) are more involved in the second

conclusion. The theorem can be applied to branched coverings which are also lamination maps. For the definitions of invariant laminations and lamination maps, and their purpose, see the extended introduction of [R2].

2.2. The Fixed Holed-Sphere Theorem. *Let $g : \bar{C} \rightarrow \bar{C}$ be an orientation-preserving critically finite degree two branched covering, with at least one non-degenerate Levy cycle, and no degenerate Levy cycle. In a neighbourhood of any periodic critical point in $X = X(g)$ fixed by g^p , let there exist a local topological conjugacy to $z \mapsto z^n$ for suitable n .*

Then there is an open subset Y of $\bar{C} \setminus X$, and an integer m , such that $\bar{C} \setminus Y$ has $\ell \geq 2$ components, each intersecting X , $g^m|_{\bar{Y}}$ is a homeomorphism, Y and $g^m Y$ are isotopic in $\bar{C} \setminus X$, with g^m cyclically permuting the boundary components, and one of the following holds.

Expanding Fixed Set Case: $m = 1$ and $\bar{Y} \subset gY$.

Nonrational Adjacent Case: $m\ell$ is less than the period of any point in X and $\partial Y = \partial_1 \cup \partial_2$, where ∂_1 and ∂_2 have the following properties. Each ∂_i is a finite disjoint union of closed arcs, two on each component of ∂Y , $\partial_1 \cap \partial_2$ is the set of endpoints of these arcs, and $\partial Y \cap g^m \bar{Y} \subset \partial_1$, $\partial g^m Y \cap \bar{Y} \subset g^m \partial_2$. Each component of $Y \setminus g^m Y$ has one component of intersection with each of ∂_1 and $g^m \partial_1$, and 2 components of intersection with ∂_2 . Each component of $g^m Y \setminus Y$ has one component of intersection with each of ∂_2 and $g^m \partial_2$, and 2 components of intersection with $g^m \partial_1$.

See Diagram 3 for what Y looks like in the nonrational adjacent case.

Diagram 3

2.3. Different Cases of, and Additions to, the Fixed Holed-Sphere Theorem.

The different cases of the Fixed Holed-Sphere Theorem are dependant on the different conclusions of the Decomposition Proposition of Chapter 1. All the proofs start with the same basic idea, but then develop differently. Let f be the branched covering of the Decomposition Proposition which is equivalent to g via a homotopy $\{f_t\}$, such that f has a complete Levy set Γ , fixed loop or gap P . The Expanding Fixed Set Case of the Fixed Holed-Sphere Theorem holds when either of the first two cases of the Decomposition Proposition hold, that is, the Rational Critical Branched Covering Case, or the Separating Periodic Set Case. The Nonrational Adjacent Case of the Fixed Holed Sphere Theorem holds when the critical branched covering of f is nonrational and the Adjacent Periodic Set Case of the Decomposition Theorem holds.

Some unfortunately complicated variations on the Fixed Holed Sphere Theorem will be needed later, when we come to apply this result to lamination maps. We state these now.

Variation in the Fixed Set Case. *If the condition $\bar{Y} \subset gY$ is weakened to $\bar{Y} \subset g^n Y$ for some n , then, given $M > 0$, we can ensure that, for any arc γ such that $g^k \gamma$ is of length $\leq M$, for all sufficiently large k , then $\gamma \subset Y$.*

Variations in the Nonrational Adjacent Case.

1. *Given an integer p , we can ensure that $g^{im}(g^m Y \setminus Y) \cap Y = \emptyset$ for $0 \leq i \leq p$.*
2. *Given $M'' > 0$, Y can be chosen so that the length of any path in $Y \setminus g^m Y$ joining ∂_2 and $g^m \partial_2$ is $> M''$.*
3. *If g is a lamination map preserving a lamination L , and I_i ($1 \leq i \leq r$) is any finite collection of S^1 -segments between noncompact leaves in gap polygons, then $\partial_1 \cup \partial_2$ can be chosen to be a union of leaf segments and S^1 -segments not including any I_i .*

2.4. Outline of proof of the Fixed Holed-Sphere Theorem.

We can assume that the homotopy f_t from g to f satisfies $f_t = g$ on a neighbourhood U of the forward orbits of periodic critical points. Then as in the semiconjugacy proof of Chapter 4 of [R2], we can find homeomorphisms Ψ_t ($t \in [0, 1]$) with

$$\Psi_0 = \text{identity},$$

$$\Psi_t = \text{identity on } U \cup X,$$

$$f_t \circ \Psi_t = g.$$

Then $f \circ \Psi_1 = g$, and $\Psi_1(g^{-1}U) = f^{-1}U$. Then define Ψ_{t+n} ($n \geq 1$) by

$$f^n \circ \Psi_{t+n} = \Psi_t \circ g^n,$$

$$\Psi_{t+n} = \text{identity on } U \cup X.$$

Then $\Psi_{t+n+1} = \Psi_{t+n}$ on $g^{-n}U$ for $t \in [0, 1]$. We also have

$$f \circ \Psi_{n+1} = \Psi_n \circ g,$$

since $f^n \circ f \circ \Psi_{n+1} = f^{n+1} \circ \Psi_{n+1} = g^{n+1} = g^n \circ g = f^n \circ \Psi_n \circ g$. Our set Y in the Rational Critical Branched Covering and Periodic Separating Set Cases will be obtained from sets of the form $\Psi_n(Z)$, where Z is a neighbourhood of P .

2.5. Proof of the Fixed Holed-Sphere Theorem: Rational Critical Branched Covering Case.

This is the most satisfactory case to deal with. We define a continuous semimetric d on $\bar{\mathbf{C}}$ with the following properties.

1. There is $\delta_0 > 0$ such that, given $\delta_1 > 0$, there are C and $\lambda > 1$ such that: if $d(f^i x, f^i y) < \delta_0$ for $0 \leq i < n$ and $d(f^i x, X) > \delta_1$ for $0 \leq i < n$ then $d(f^n x, f^n y) \geq C\lambda^n d(x, y)$.

Note, however, that we might have $d(x, y) = 0$ and $d(fx, fy) > 0$.

2. If a set $A = \{x : d(x, x_0) = 0\}$ is noncontractible, then it is mapped by f to a set of the form $\{y : d(y, y_0) = 0\} = B$. If A is contractible then ∂A is mapped into ∂B for some such set B .

3. $d(x, y) = 0$ for all $x, y \in P$. If $\varepsilon > 0$ is sufficiently small, then $\{x : d(x, P) < \varepsilon\}$ can be homotoped in $\bar{\mathbf{C}} \setminus X$ into an arbitrarily small neighbourhood of P .

4. $\{y : d(x, y) < \varepsilon\}$ is always connected.

Suppose this semimetric has been defined. Then as in Chapter 4 of [R2] (that is, using property 1 above) we obtain from the equation $f^n \circ \Psi_{t+n} = \Psi_t \circ g^n$ that

$$d(\Psi_{n+1}(x), \Psi_n(x)) \leq C_1 \lambda^{-n},$$

for all x and for some $C_1 > 0$ and $\lambda > 1$. We use the fact that $\Psi_n = \Psi_{n+1}$ on $g^{-n}U$. Similarly, if γ is such that $g^n \gamma$ always has length $\leq M$, then $\lim_{n \rightarrow \infty} \text{diameter } \Psi_n(\gamma) = 0$. (See the variations on the Fixed Holed Sphere Theorem.) Then we can find $\delta_2 > 0$, N and $\mu > 1$ such that, $d(X, P) > 2\delta_2$, and if

$$d_N(x, y) = \sum_{i=0}^N d(f^i x, f^i y),$$

and $d_N(x, y) < \delta_2$, $d(x, X) > \delta_0$, then $d_N(fx, fy) \geq \mu d_N(x, y)$. Then for $0 < \varepsilon < \delta_0$, consider

$$Y' = \{x : \lim_{n \rightarrow \infty} d_N(\Psi_n(x), P) < \varepsilon\},$$

which is automatically open. Then since, if $Z = \{z : d_N(z, P) < \varepsilon\}$, $\bar{Z} \subset fZ$, and $f|Z$ is a homeomorphism for ε small enough, $\bar{Y}' \subset gY'$ and $g|Y'$ is a homeomorphism. Let

$$Y = \text{Interior}(Y' \cup (\text{components of } \bar{\mathbf{C}} \setminus Y' \text{ disjoint from } X)).$$

Then Y has the required properties.

We can, in fact, do much more than construct d : we can construct a semiconjugacy π between g and $[f]$ on a certain subset of $\bar{\mathbf{C}}$, with dense image under π , where $[f]$ is as follows. Let $\sim$ denote the equivalence relation $x_1 \sim x_2$ if and only if $d(x_1, x_2) = 0$. Let $\pi : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}/\sim$ be the quotient map. Let $[f]([x]) = [f(x)]$ if $f([x]) = [f(x)]$ (where $[x]$ is the equivalence class of x), and in general $[f]([x]) = [f(x')]$ for any $x' \in \partial[x]$. By property 2 of d , $[f]$ is well-defined. Then $\pi \circ f = [f] \circ \pi$ on a subset of $\bar{\mathbf{C}}$. It follows that $\pi \circ \Psi_n \circ g = [f] \circ \pi \circ \Psi_{n+1}$ on a subset A_n of $\bar{\mathbf{C}}$. Then we can show that $\pi \circ \Psi_n$ converges to a map $[\Psi]$ satisfying $[\Psi] \circ g = [f] \circ [\Psi]$ on $\limsup_{n \rightarrow \infty} A_n$.

2.6. Construction of the semi-metric.

Recall that the critical gap (see 1.8) G_0 is periodic under f_* , and that the critical branched covering is equivalent to a rational map, since we are in the Rational Critical Branched Covering case. Let us call the critical branched covering h_0 . The critical points of h_0 may not be periodic. But the forward orbit of any periodic critical point is in G_0 . We originally chose h_0 (in 1.8) with $h_0^{-1}G_0 \subset G_0$, $\partial G_0 \subset \partial h_0^{-1}G_0$. This is consistent with choosing h_0 so that any periodic critical point is locally conjugate to $z \mapsto z^p$ for suitable p , as we now do. We can also assume that $h_0 = f^n|_{f^{-n}G_0 \cap G_0}$, if G_0 has period n . Now by the Semiconjugacy Proposition of 4.1 of [R2], if h_1 is the rational map equivalent to h_0 , there is a semiconjugacy $\psi : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ with $\psi \circ h_0 = h_1 \circ \psi$, and ψ must collapse each component of $\bar{\mathbf{C}} \setminus G_0$ to a point. (We do not really need the Semiconjugacy Proposition. We could just be more careful in the choice of h_0 , so that it would be clear that h_1 existed.) Let d_s denote the spherical metric on $\bar{\mathbf{C}}$, and define

$$d_{G_0}(x, y) = d_s(\psi x, \psi y).$$

Then d_{G_0} satisfies property 1 required for d .

Now let G be any component of $\cup_{n \geq 0} f^{-n}G_0$ which is not contained in a disc D such that all components of $f^{-m}D$ are discs for all $m \geq 0$. Let $\mathcal{G}_n$ be the set of those components G for which n is the least integer ≥ 0 with $f^n G = G_0$. Then for $G \in \mathcal{G}_n$, define

$$d_G(x, y) = (1/3^n)d_{G_0}(f^n x, f^n y).$$

Note that, from the definition of $\mathcal{G}_n$, $f^n : G \rightarrow G_0$ is a homeomorphism for $G \in \mathcal{G}_n$.

Now define d as follows. For any $x, y \in \bar{\mathbf{C}}$, choose a path $\alpha : [0, 1] \rightarrow \bar{\mathbf{C}}$ with $\alpha(0) = x$, $\alpha(1) = y$, and such that α has at most one component of intersection with any $G \in \cup_{n \geq 0} \mathcal{G}_n$. Then write

$$\alpha^{-1}(G) = (s_G, t_G),$$

if $\alpha^{-1}(G) \neq \phi$. Then define

$$d(x, y) = \sum_{\alpha^{-1}(G) \neq \phi} d_G(\alpha(s_G), \alpha(t_G)).$$

Then d is well-defined and finite, since $\#(\mathcal{G}_n) \leq 2^n$, so that

$$\sum_{n=0}^{\infty} \sum_{G \in \mathcal{G}_n} (1/3^n) < \infty.$$

It is then easy to check that d has the required properties.

2.7. Proof of the Fixed Holed-Sphere Theorem: A preliminary lemma in the Periodic Separating Set Case.

In this case, Q exists of period ℓ under f_* , separating P from f_*G_0 . See 1.14 for the following. There is a local inverse T of f^ℓ defined on $P \cup R$, where R is the set bounded by f_*G_0 and P , with $TP = P$ and $TR \subset R$. There is a local inverse $S \neq T$ of f^ℓ defined on a neighbourhood of Q with $SQ = Q$, such that S either reverses orientation on Q , if Q is a loop, or cyclically permutes the boundary components of Q (if Q is a gap) and S^k fixes them, for some $k \geq 3$. Thus, TQ separates P and Q , and $T(\partial R \cap \partial f_*G_0)$ - in the boundary of a component G_1 of $\cup_{n>0} f^{-n}G_0$ - separates Q and TQ .

Lemma. *There are G_i, G'_i ($i \geq 1$) between Q and TQ , such that*

- a) G'_i separates G_j from TQ for all i, j ,
- b) G_i separates G_j from Q , and G'_i separates G'_j from TQ , if $i > j$,
- c) there is N_1 such that

$$G_i, G'_i \subset \bigcup_{j \leq N_1 + i\ell k} f^{-j}G_0.$$

See Diagram 4.

Proof: $S^p G_1$ separates Q and f_*G_0 for some $1 \leq p < k$. Then we can take $G_i = S^{k(i-1)}G_1$ and $G'_i = TS^{p+k(i-1)}G_1$. Thus, G_i, G'_i are in the disjoint sets SR, TR respectively, and have all the required properties.

2.8. Proof of the Fixed holed Sphere Theorem in the Periodic Separating Set Case.

Let $Z_{0,i}, Z_1, Z_{2,i}, Z_3, Z_{4,i}, Z_{5,i}, Z_6$ be the open sets containing P and bounded by the images under f^j ($0 \leq j < \ell$) of $G'_i, TQ, TG_i, T^2Q, T^2G_i, T^2G'_i, T^3Q$ respectively. Then

$$\bar{Z}_6 \subset Z_{5,i+1} \subset Z_{5,i} \subset \bar{Z}_{5,i} \subset Z_{4,i} \subset \bar{Z}_{4,i} \subset Z_{4,i+1} \subset Z_3 \subset \bar{Z}_3 \subset Z_{2,i} \subset \bar{Z}_{2,i}$$

Diagram 4

$$\subset Z_{2,i+1} \subset Z_1 \subset \bar{Z}_1 \subset Z_{0,i+1} \subset \bar{Z}_{0,i+1} \subset Z_{0,i} \quad (1)$$

We take

$$Y_{j,i,n} = \Psi_n^{-1} Z_{j,i},$$

if $j = 0, 2, 4$ or 5 , and

$$Y_j = \{z : \overline{\{\Psi_n(z) : n \geq N\}} \subset Z_j \text{ for some } N\}$$

if $j = 1, 3$ or 6 . We shall find N_2 such that, if $n, n' \geq N_2 + i\ell k$, for any i ,

$$Y_6 \subset \bar{Y}_6 \subset Y_{5,i,n} \subset Y_{4,i,n} \subset \bar{Y}_{4,i,n} \subset Y_1 \subset \bar{Y}_1 \subset Y_{0,i,n'}. \quad (2)$$

Then we take Y to be the union of the component of $\text{Interior}(Y_6)$ which is nontrivial in $\bar{\mathbf{C}} \setminus X$, together with all complementary components which do not intersect X . Then $g^{2\ell}Y$ is the component of $\text{Interior}(Y_1)$ which is nontrivial in $\bar{\mathbf{C}} \setminus X$, together with all complementary components which do not intersect X . Since $f^{2\ell} : \bar{Z}_{5,i} \rightarrow \bar{Z}_{0,i}$ is a homeomorphism, so is $g^{2\ell} : \bar{Y}_{5,i,n+2\ell} \rightarrow \bar{Y}_{0,i,n}$, and if (2) holds,

$$\bar{Y} \subset \bar{Y}_{5,i,n+2\ell} \subset Y_{4,j,n+2\ell} \subset g^{2\ell}Y. \quad (3)$$

This is enough to show that Y is nonempty and homotopic to P in $\bar{\mathbf{C}} \setminus X$. This is the set Y that we need for the variation 2.3 on the Fixed Holed-Sphere Theorem, rather than the Fixed Holed-Sphere Theorem itself, for which we would need to replace Y by a union of $\cap_{i=0}^{2\ell} g^i Y$ and appropriate complementary components.

It remains to show (2). For this, it suffices to show that, for $n \geq N_2 + i\ell k$, and $n \leq n' \leq n + 2\ell k$,

$$Y_{5,i+2,n'} \subset Y_{5,i,n} \subset Y_{4,i,n} \subset Y_{4,i+2,n'}, \quad Y_{0,i+2,n'} \subset Y_{0,i,n}. \quad (4)$$

For then,

$$\overline{\{\Psi_{n'}(z) : n' \geq n\}} \subset \bar{Z}_3 \subset Z_1$$

whenever $z \in Y_{4,i,n}$, and so $\bar{Y} \subset Y_1$. Also,

$$\bigcap_{n \geq N} \bigcap_i Y_{0,i,n} = \bigcap_{n \geq N_2} \bigcap_i Y_{0,i,n}$$

if $N \geq N_2$. So $Y_1 \subset \bigcap_i Y_{0,i,n}$ if $n \geq N_2$, and $\bar{Y}_1 \subset Y_{0,i,n}$. To show (4), it suffices to find N_3 such that, for $n \geq N_3 + i\ell k$, and any x , $\{\Psi_{n+t}x : 0 \leq t \leq 2\ell k\}$ crosses at most one set of the form G'_j or T^2G_j or $T^2G'_j$, for $j \leq i + 2$.

To complete the proof of the Fixed Holed-Sphere Theorem and its variations in this case, assuming (4) holds, we have to show that if γ is any arc such that for all sufficiently large n , has $g^n\gamma$ arc length $\leq M$, and $\gamma \cap Y \neq \emptyset$, then $\gamma \subset Y$. It suffices to show this for Y_1 replacing Y , since $g^{2\ell}Y$ is obtained by adding some discs to Y_1 . So fix $x \in \gamma$, and suppose $\{\Psi_n(x) : n \geq N\} \subset Z_1$ for some N . We need to show there is N' such that $\Psi_n(\gamma) \subset Z_1$ for all $n \geq N'$. But $\{\Psi_n(x) : n \geq N\} \subset Z_{2,i}$ for some i . So it suffices to find $N_3 \geq N$ such that $\Psi_n(\gamma)$ crosses at most one set TG_s for $s \leq j$ and $n \geq N_3 + j\ell k$. For then $\Psi_n(\gamma) \subset Z_{2,i+2} \subset Z_1$ for $n \geq N_3 + i\ell k$, and $\gamma \subset Y_1$, as required.

Now $\Psi_n(\gamma)$ and $\{\Psi_{n+t}x : 0 \leq t \leq \ell k\}$ are both components of the inverse image under f^n of arcs of bounded length. So the proof of both (4) and the necessary fact about $\Psi_n(\gamma)$, and hence the proof of the theorem, is completed by the following lemma.

2.9. Lemma.

Given $M > 0$, there is N_4 such that the following holds. If α is any arc of length $\leq M$, then, for $n \geq N_4$, any component of $f^{-n}\alpha$ intersects at most one component of $\bar{\mathcal{C}} \setminus \bigcup_{i \geq 0} f_^i G_0$.*

Proof: Recall that Γ is a complete Levy set for f , and let $\Gamma_1 = f^{-r}\Gamma$ for some suitable r , such that a nonperiodic loop of Γ_1 separates Q_1 and Q_2 , whenever these are disjoint closed sets such that Q_i is either a periodic loop in Γ or the closure of a periodic gap for Γ which is always mapped homeomorphically by f^n . This is clearly possible, because of the definition of complete Levy set (see Chapter 1) and because if G_0 is periodic, only one boundary component is periodic. Then it is clear that there exists L for which the following is true. For any α of length $\leq M$, there exist $L' = L'(\alpha) \leq L$ and α_i ($0 \leq i \leq L'$) with $\alpha = \bigcup_{i=1}^{L'} \alpha_i$, such that the arc α_i is homotopic in $\bar{\mathcal{C}} \setminus X$, via a homotopy fixing endpoints, to one with at most one intersection with a loop in Γ_1 . Then there is N_4 (independent of α) such that for $n \geq N_4$, any component of $g^{-n}\alpha$ intersects at most one element of Γ_1 , since it intersects at most L' elements of $g^{-n}\Gamma_1$. Then each one can intersect at most one component of $\bar{\mathcal{C}} \setminus \bigcup_{i \geq 0} f_*^i G_0$, since the boundary of each $f_*^i G_0$ is contained in $\Gamma \subset \Gamma_1$.

2.10. Proof of the Fixed Holed-Sphere Theorem: a simpler analogue of the Nonrational Adjacent Case.

Let $g : \mathbf{R}^2/\mathbf{Z}^2 \rightarrow \mathbf{R}^2/\mathbf{Z}^2$ be a covering of the two-torus homotopic to an endomorphism given by a matrix A in $GL(2, \mathbf{Z})$, where A has determinant 2 and eigenvalues λ_u, λ_s with $|\lambda_s| < 1 < 2 < |\lambda_u|$. Let v_u, v_s be the corresponding eigenvectors. Let $\bar{g}$ denote a (well-defined) lift of the multivalued function g^{-1} to $\mathbf{R}^2$ (so that notation agrees with what we shall do later). Then let

$$V = \{xv_u + yv_s : |x| < M'\},$$

$$W = \{xv_u + yv_s : |y| < M'\},$$

for some suitable $M' > 0$. We have $AW \subset W$, because $A(xv_u + yv_s) = x\lambda_u v_u + y\lambda_s v_s$, and similarly $A^{-1}V \subset V$. In fact, the distance between the boundaries of W and AW increases with M' , and similarly with $V, A^{-1}V$. But there is M such that $\text{distance}(\bar{g}x, A^{-1}x) \leq M$ for all x . So if M' is sufficiently large given M , we can ensure that $\bar{g}V \subset V$, and similarly $W \subset \bar{g}W$.

Now we consider the set $\bar{g}^n(V \cap W) = Y_n$, so that $Y_{n+1} = \bar{g}Y_n$. Let

$$B_n = \{z \in \mathbf{Z}^2 : x, x+z \in Y_n \cup Y_{n+1} \text{ for some } x\}.$$

Then $B_n = A^{-n}B_0 \cap \mathbf{Z}^2$. We claim that $\bigcap_{n=0}^{\infty} A^n \mathbf{Z}^2 = \{0\}$. For A is an automorphism of the subgroup $\bigcap_{n=0}^{\infty} A^n \mathbf{Z}^2$, which must then be either $\{(0,0)\}$ or rank two, since no nontrivial rationally defined subspace is invariant under A . But A has determinant 2, so this subgroup must indeed be $\{(0,0)\}$. (I would like to thank my colleagues for helping me clarify this rather trivial point.) So, since B_0 is finite, $B_n = \{(0,0)\}$ for all sufficiently large n . So for sufficiently large n , $Y_n \cup Y_{n+1}$ projects homeomorphically to $\mathbf{R}^2/\mathbf{Z}^2$, and g maps the projection of Y_{n+1} homeomorphically to the projection of Y_n . Moreover, if Y denotes the projection of Y_{n+1} , we can write $\partial Y = \partial_1 \cup \partial_2$, where each ∂_j is the union of two disjoint arcs, and $g\partial_1 \cap Y = \phi$, $gY \cap \partial_2 = \phi$. This is about as far as we can take the arguments of [Fr]. We now want to generalise this argument, to prove the Nonrational Adjacent case of the Fixed Holed-Sphere Theorem.

2.11. Proof of the Fixed Holed-Sphere Theorem in the Nonrational Adjacent Case: lifting the branched covering inverse to the unit disc.

We are working under the following assumptions. The map $g : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ is a degree two critically finite orientation-preserving branched covering, with nonrational critical branched covering and an adjacent periodic set. (See the Decomposition Proposition.) Thus, there is a homotopy $\{f_t : t \in [0,1]\}$ through critically finite branched coverings with $f_0 = g$, $f_1 = f$, $X(f_t) = X(f) = X$ for all $t \in [0,1]$, and f has the following properties.

The set Q . There is a set Q such that $f^m Q = Q$, for some $m \geq 1$. Either Q is a loop, in which case f^m reverses orientation on Q and we write $\ell = 2$, or Q

is a ℓ -holed sphere (for some $\ell \geq 3$). In the latter case, f^m cyclically permutes the components of ∂Q , and f^q fixes the boundary components, where $q = m\ell$.

The sets $G, \hat{G}$. Either way, there is a set G adjacent to Q with the following properties. The sets $f^i G$ ($0 \leq i \leq q$) are disjoint for $i < q$, and adjacent to Q when $m \mid i$, with $G \subset f^q G$, $\partial f^q G \subset \partial G$. Also, G has a holed torus as a double covering. There is a holed plane $\hat{G}$ covering this torus, and $f^{-q} \mid G$ lifts to $\hat{f} : \hat{G} \rightarrow \hat{G}$ having the following properties. The map $\hat{f}$ is injective, but not surjective, and there are a C^1 $\varphi : \hat{G} \rightarrow \mathbf{R}^2$ and $A \in GL(2, \mathbf{Z})$, with determinant 2, such that

$$\varphi = A \circ \varphi \circ Acc\hat{f}.$$

Moreover, φ conjugates the action of the covering group of the torus to the action of $\mathbf{Z}^2$ on $\mathbf{R}^2$ by translation. The eigenvalues λ_u, λ_s of A satisfy $|\lambda_s| < 1 < 2 < |\lambda_u|$. We fix corresponding eigenvectors v_u, v_s .

Lifting Inverses, and Paths Representing Points in the Universal Cover. Now we recall how to lift f_t^{-n} ($n \geq 1$) to the universal cover $\mathring{D}$ of $\bar{\mathbf{C}} \setminus X$ (This lifting of the inverses of critically finite branched coverings is fundamental in [T].) We fix a basepoint x_1 in $\bar{\mathbf{C}} \setminus X$ - which, for convenience, we take in Q . Fix n , and fix x'_1 with $f_1^n x'_1 = x_1$, hence fixing a path $\{x'_t\}$ with $f_t^n x'_t = x_1$. We also fix a path α_1 from x_1 to x'_1 . Then α_t is the path α_1 followed by $\{x'_u\}$ from x'_1 to x'_t . Then $\mathring{D}$ can be identified with homotopy classes of paths in $\bar{\mathbf{C}} \setminus X$ based at x_1 . If β is any such path, let β_t be the component of $f_t^{-n}\beta$ based at x'_t . We write

$$\bar{f}_t[\beta] = [\alpha_t * \beta_t],$$

where $[\beta]$ is the homotopy class of β . Then $\bar{f}_t : \mathring{D} \rightarrow \mathring{D}$ is well-defined.

The group Δ . If $\bar{f}_t[\beta] = \bar{f}_t[\gamma]$, and $[\beta] \neq [\gamma]$ then $[\beta] = [\tau * \gamma] = [\tau] \cdot [\gamma]$, where τ is a closed loop in $\bar{\mathbf{C}} \setminus X$ based at x_1 (so that $[\tau] \in \pi_1(\bar{\mathbf{C}} \setminus X)$). The converse is not necessarily true. The set of τ which arises is independent of t . Similarly, if $\bar{f}_t[\beta] = [\tau] \cdot \bar{f}_t[\gamma]$ for some $\tau \in \pi_1(\bar{\mathbf{C}} \setminus X)$, then $[\beta] = [\sigma] \cdot [\gamma]$, where $[\sigma] \in \pi_1(\bar{\mathbf{C}} \setminus X)$ is again independent of t . Let Δ be the subgroup of $\pi_1(\bar{\mathbf{C}} \setminus X)$ consisting of all $[\delta]$ such that $\bar{f}^n([\delta] \cdot x) \in \pi_1(\bar{\mathbf{C}} \setminus X) \cdot \bar{f}^n(x)$ for all x , for all n . Then $\pi_1(Q), \pi_1(\hat{G}) \leq \Delta$. From the definition of $\bar{f}_t[\beta]$, if $\bar{f}_t[\beta] = [\gamma]$, there is δ with initial point at x'_t , and

$$[\alpha'_t \cup \delta] = [\gamma],$$

$$[f_t^n \delta] = [\beta].$$

2.12. The construction of V_t and W_t .

From now on, we take $\bar{f}_t$ to be a lift of f_t^{-m} , with $\bar{f} = \bar{f}_1$ and $\bar{g} = \bar{f}_0$. Let $\tilde{Q}$ be a fixed lift of Q , and let $\tilde{G}$ be a lift of G adjacent to $\tilde{Q}$. By taking $x_1, x'_1 \in Q$ and $\alpha_1 \subset Q$, we can ensure that $\bar{f}\tilde{Q} = \tilde{Q}$.

Now we are ready to construct ∂V_t by constructing the Δ -orbit ∂_t of its boundary, and then defining V_t to be the component of $\mathring{D} \setminus \partial_t$ containing $\tilde{Q}$. We define $\partial_t = \cup_{i=0}^{\ell-1} \bar{f}_t^i \partial'$ where ∂' is Δ -invariant, and is defined by its intersection with $\tilde{G}$. Since $\pi_1(\hat{G}) \leq \Delta$, it suffices to construct the projections of the components to $\hat{G}$. To do this, it suffices to define the images under φ in $\mathbf{R}^2$, provided these are disjoint from $\varphi(\partial \hat{G})$.

We take the images under φ of projections of components of ∂' to $\hat{G}$ to be

$$\{xv_u + yv_s : x = \pm M'\}.$$

The definition of W_t is similar, with v_u and v_s interchanged.

2.13. Lemma.

$\bar{f}V_1 \subset V_1$ and given $M > 0$, M' can be chosen so that $d_P(\partial V_1, \bar{f}\partial V_1) > M$, where the d_P is the Poincaré metric. Similarly, W_1 contains a component of $\bar{f}^{-1}W_1$ containing $\tilde{Q}$, and $d_P(\partial W_1, \bar{f}\partial W_1) > M$.

Proof: We represent points in $\mathring{D}$ by homotopy classes of paths in $\bar{\mathbf{C}} \setminus X$ with first endpoints at $x_0 \in Q$. We shall use the same letters to represent the lifts of the paths to the universal cover $\mathring{D}$. We consider the case of ∂V_1 - the proof for ∂W_1 is exactly similar. So first, let α be a path representing an element of V_1 , with first endpoint in Q . Thus, $\alpha \subset V_1$. We have to show $\bar{f}\alpha \subset V_1$. By Lemma 2.14 below, for all sufficiently large n , $\bar{f}^n \alpha$ has only one component of intersection with Q and none with any other component of $\bar{\mathbf{C}} \setminus \cup_{i=0}^{\ell-1} \bar{f}^{im} G$ (assuming that α has only essential intersections with components of $\bar{\mathbf{C}} \setminus \cup_{i=0}^{\ell-1} \bar{f}^{im} G$). For n divisible by ℓ (without loss of generality), $\bar{f}^n \alpha \subset Q \cup G$, and $\bar{f}^n \alpha$ does not cross $\bar{f}^n \partial_1$, because $\partial_1 = \bar{f}^{-n}(\bar{f}^n \partial_1)$. In $\tilde{G}$ itself, $\bar{f}^n \partial_1$ separates $\bar{f}^{n-1} \partial_1$ from $\tilde{Q}$. So $\bar{f}^n \alpha$ also does not cross $\bar{f}^{n-1} \partial_1$, and so $\bar{f}\alpha$ does not cross ∂_1 . Hence $\bar{f}\alpha \subset V_1$, as required.

To show $d_P(\partial V_1, \bar{f}\partial V_1) > M$, suppose not, and let α be a path of length $\leq M$ between the boundary components. Again by 2.14, for some n and M_1 depending only on M , $\bar{f}^n \alpha$ is a path of length $\leq M_1$ which may have one component of intersection with Q , but intersects no other component of $\bar{\mathbf{C}} \setminus \cup_{i=0}^{\ell-1} \bar{f}^{im} G$ (again, assuming only essential intersections for α). Suppose without loss of generality that part of $\bar{f}^n \alpha$ with endpoint lifting to $\bar{f}^n \partial V_1$ lies in G , and hence lifts to $\hat{G}$, with endpoint in $\varphi(\hat{f}^n \partial')$. But because $\Delta \cap \pi_1(G) = \pi_1(\hat{G})$, $\bar{f}^n \alpha$ cannot intersect Q in just one component. So we can choose n so that $\bar{f}^n \alpha \subset G$,

and then obtain a bound (in terms of M) on the distance between components of $\bar{f}^n \partial_1 \cap \tilde{G}$ and $\bar{f}^{n+1} \partial_1 \cap \tilde{G}$, which can be contradicted by choice of M' .

2.14. Lemma. *Let M be given. Then there is r such that, if $\gamma : [0, 1] \rightarrow \bar{\mathcal{C}} \setminus X$ is any path of length $\leq M$, the following holds after homotopy of γ in $\bar{\mathcal{C}} \setminus X$ fixing the endpoints. For $n \geq r$, a lift of γ under f^n has at most one component of intersection with Q , and none with any other component of $\bar{\mathcal{C}} \setminus \cup_{i=0}^{\ell-1} f^{im} G$.*

Proof: We can homotope γ , fixing endpoints, to have $m_0 < \infty$ intersections with $\cup_{i=0}^{\ell-1} f^{im} \partial G$, all essential in $\bar{\mathcal{C}} \setminus X$. Let m_n be the maximal number of intersections for a lift of γ under f^n . Then since the intersections for a lift under f^n are inverse images under f^n of the intersections for γ , $\{m_n\}$ is a decreasing sequence, and there is r_1 such that, for $n \geq r_1$, the intersections are only with components of ∂Q . Now if γ' is any path of length $\leq M_2$ with endpoints in ∂Q but otherwise lying in some $f^{im} G$, there is r_2 (depending only on M_2) such that, for $n \geq r_2$, no lift of γ' under f^n has both endpoints in ∂Q . Applying this repeatedly, we obtain the required r .

2.15. Properties of V_t, W_t .

Note that there is M such that, for $0 \leq i \leq \ell$,

$$d_P(\bar{f}^i x, \bar{f}_t^i x) < M \quad (1).$$

Then if M' is sufficiently large given M , we have the following corollary of 2.13.

Corollary.

$\bar{f}_t^{n+1} \bar{V}_t \subset \bar{f}_t^n V_t$, $\bar{f}_t^n \bar{W}_t \subset \bar{f}_t^{n+1} W_t$, $\partial \bar{f}_t^n V_t = \bar{f}_t^n \partial V_t$, and $\partial \bar{f}_t^n W_t = \bar{f}_t^n \partial W$ for all $t \in [0, 1]$ and $n \geq 0$.

Proof: By 2.13 and (1) above, $\bar{f}_t \partial V_t \subset \bar{V}_t$. The last two statements follow from the fact that $\partial(\Delta \cdot \bar{f}_t^n V_t) = \Delta \cdot \bar{f}_t^n \partial V_t$, and similarly for W_t . This is easily proved by induction: the point is that $\bar{f}_t$ is a covering from $\Delta \cdot \bar{f}_t^n \bar{V}_t$ onto $\Delta \cdot \bar{f}_t^{n+1} \bar{V}_t$, and similarly for W_t . Hence $\bar{f}_t^n V_t \subset V_t$ and by induction $\bar{f}_t^{n+1} V_t \subset \bar{f}_t^n V_t$. The proof that $\bar{f}_t^n W \subset \bar{f}_t^{n+1} W_t$ is similar, and actually slightly easier.

2.16. Proof of the Fixed Holed-Sphere Theorem in the Nonrational Adjacent Case.

Note that $V_1 \cap W_1$ has the following property: for some integer N , $\bar{f}^n(\bar{V}_1 \cap \bar{W}_1)$ projects to a compact subset of $\mathring{D}/\pi_1(Q)$ for $n \geq N$. This is simply because, for some N the set

$$A^{-n}(\{xv_u + yv_s : |x| \leq M', |y| \leq M'\})$$

does not intersect $\mathbf{Z}^2 \setminus \{(0,0)\}$ for $n \geq N$. We claim that $\bar{f}_t^n(\bar{V}_t \cap \bar{W}_t)$ also projects to a compact subset for all t . For

$$\bar{f}_t^n(\bar{V}_t \cap \bar{W}_t) \subset \bar{f}^n(\bar{V}_1 \cap \bar{W}_1) \cup \bigcup_{0 \leq i < \ell, t \leq u \leq 1} \bar{f}_u^i(\psi_u \bar{f}^n \partial(V_1 \cap W_1)).$$

Here, ψ_u is a lift of the homeomorphism $f_u^{-n} \circ f^n$ to $\mathring{D}/\pi_1(Q)$. So the claim is proved. So, in particular, $\bar{g}^N(V_0 \cap W_0) = [\tau] \cdot \bar{g}^N(V_0 \cap W_0)$ for $[\tau]$ lying in only finitely many cosets of $\pi_1(Q)$. Then 2.14 implies that by enlarging N , we can assume this is true only for $[\tau] \in \pi_1(Q)$. Take Y to be the projection of $\bar{g}^N(V_0 \cap W_0)$ to $\bar{\mathbf{C}} \setminus X$. Let ∂_1 be the projection of $\bar{g}^N(\partial V_0 \cap \partial(V_0 \cap W_0))$, and ∂_2 the projection of $\bar{g}^N(\partial W_0 \cap \partial(V_0 \cap W_0))$. If we isotope ∂' (see 2.12) a distance $\leq M$ in a Δ -invariant way, and redefine V_t using the new definition of ∂' , then V_t still has the properties claimed above, and similarly for W_t . So we can assume that each component of ∂V_0 , or $\bar{g}\partial V_0$ intersects each component of ∂W_0 at most once, by choice of ∂' and the corresponding set for ∂W_0 . Then Y has all the properties stated in the Fixed Holed-Sphere Theorem.

2.17. Proof of the Variations in the Nonrational Adjacent Case.

For any $p > 0$, the set $\cup_{i=0}^p \bar{g}^{N-i}(V_0 \cap W_0)$ intersects its image under $[\tau]$ only for $[\tau] \in \pi_1(Q)$ if N is large enough. We can also make the distance in $\bar{g}W_0$ between ∂W_0 and $\bar{g}\partial W_0$ arbitrarily large by choice of M' - and then the distance in $\bar{g}^{n+1}W_0$ between $\bar{g}^nW_0$ and $\bar{g}^{n+1}\partial W_0$ is even larger, for any $n > 0$. If g is a lamination map, we can choose ∂' (and the analogue used to define ∂W_0) to be a union of lifts of S^1 -segments and leaf segments, and by tracing round the boundaries of gaps, can avoid the lifts of the forward orbits of the S^1 -segments I_i (see variation 3 of 2.3). Then Y has all the required properties.

Chapter 3. One-dimensional Topological Dynamics.

3.1. Definitions.

Let T be a locally finite graph. If $x \in T$, $U \subset T$ is a *1-sided neighbourhood* of x if $U \setminus \{x\}$ is connected, nonempty, contains no vertex of T , and $x \in U$. (To fit in with what we need to prove, this definition and the following one may be a bit unorthodox.) Let $h : T \rightarrow T$ be continuous with $hx = x$. Then we define $W^u(x, h, U)$ by

$$W^u(x, h, U) = \{x\},$$

if arbitrarily small 1-sided neighbourhoods V of x in U satisfy $hV \setminus U \neq \emptyset$, and

$$W^u(x, h, U) = \bigcap_{n \geq 0} \left\{ \bigcup_{n \geq 0} h^n V : V \text{ is a 1-sided neighbourhood of } x, V \subset U \right\}$$

otherwise. Thus $W^u(x, h, U_1) = W^u(x, h, U_2)$ if U_1, U_2 are 1-sided neighbourhoods with $U_1 \cap U_2 \neq \emptyset$.

3.2.

The main purpose of this chapter is to prove the following.

The 1-sided Neighbourhood Theorem.

Let $Y \subset \bar{\mathbf{C}}$ be an ℓ -holed sphere with boundary ($\ell \geq 2$). Let $T \subset Y$ be a graph such that $T \hookrightarrow Y$ is a homotopy equivalence. Let $\chi : Y \rightarrow Y$ be an orientation-preserving homeomorphism which cyclically permutes the components of ∂Y . Let $h : T \rightarrow T$ be continuous and homotopic to $\chi|_T$ as a map from T to Y .

Then there is a graph $T_1 \subset T$ with no free vertices such that $\bar{\mathbf{C}} \setminus T_1$ has k components (for some $2 \leq k \leq \ell$), $h|_{T_1}$ is homotopic as a map from T_1 to T to some $h_0 : T_1 \rightarrow T_1$ which cyclically permutes the boundaries of the components of $\bar{\mathbf{C}} \setminus T_1$ up to homotopy, and

$$T_1 \subset \bigcup \{W^u(x, h^k, U) : h^k x = x, x \in T_1, U \text{ is a 1-sided neighbourhood of } x \text{ in } T_1\}$$

3.3.

The main tool in the proof of the 1-sided Neighbourhood Theorem is the Pseudo-Anosov Proposition below. Some of the same ideas occurred in the proof of the Tuning Proposition in Chapter 7 of [R2], and, as there, we rely heavily on Thurston's description of surface homeomorphisms up to isotopy, as described in [F-L-P].

Pseudo-Anosov Proposition.

Let Y_1 be a k -holed sphere ($k \geq 2$). Let $\chi : Y_1 \rightarrow Y_1$ be a homeomorphism which cyclically permutes the boundary components of Y_1 and does not leave isotopically invariant any nontrivial nonboundary disjoint loop set in Y_1 . Suppose also that χ fixes a point x_0 in Y_1 . Then χ is isotopic, via an isotopy fixing x_0 , to $\chi_0 : Y_1 \rightarrow Y_1$ satisfying one of the following.

1. $\chi_0^k = \text{identity}$.
2. χ_0 fixes a disc D round x_0 , and $\chi_0 : Y_2 \rightarrow Y_2$ is pseudo-Anosov, where $Y_2 = Y_1 \setminus D$. In particular, a measured foliation on Y_2 with singularities is preserved by χ_0 , and this foliation has singular leaf ℓ with endpoint at $x_1 \in \partial Y_1 \subset \partial Y_2$ such that x_1, ℓ are preserved by χ_0^k . Moreover, there is a length measurement defined on ℓ , and $\lambda > 1$, such that χ_0^k multiplies length on ℓ by λ , and the length of any segment of ℓ is proportional to the diameter of a lift to $\tilde{Y}_2$, where $\tilde{Y}_2$ is the universal cover of Y_2 .

Proof: To follow the outline of [F-L-P] more closely, we assume (as we may do) that χ preserves D . The Teichmuller space of $Y_2 = Y_1 \setminus D$ is then the space of (non-complete) hyperbolic structures on Y_2 for which all the boundary components are geodesics of length 1. This space is homeomorphic to the open unit ball in $\mathbf{R}^{2k-4}$ (or a single point if $k = 2$) and its Thurston compactification is homeomorphic to the closed unit ball in $\mathbf{R}^{2k-4}$. Exactly as in [F-L-P], g induces a homeomorphism of this Thurston compactification (the only difference is that, here, χ does not fix boundary components) and must have a fixed point. Since no simple disjoint nontrivial nonboundary loop set in Y_1 is left invariant by χ up to isotopy, and χ cyclically permutes the components of ∂Y_1 , there is similarly no such loop set in Y_2 . So there is χ_0 isotopic to χ such that either

1. χ_0 is an isometry, for some hyperbolic structure on Y_2 , or
2. χ_0 is pseudo-Anosov, in a sense made more precise below (and see [F-L-P]).

Case 1. We claim that $\chi_0^k = \text{identity}$ on Y_2 , and we can then extend χ_0 to Y_1 with $\chi_0^k = \text{identity}$ on Y_1 . For let γ be a geodesic arc of minimal length joining ∂D to ∂Y_1 . Then γ is disjoint from all other such arcs. It follows that $\chi_0^i \gamma$ ($0 \leq i < k$) are disjoint, and $\chi_0^k \gamma = \gamma$. Since $Y_1 \setminus (D \cup \bigcup_{i \geq 0} \chi_0^i \gamma)$ is a disc, $\chi_0^k = \text{identity}$ on Y_2 .

Case 2. We are only really interested in the unstable foliation preserved by χ_0 . This singular foliation has finitely many singularities, at least one on each component of ∂Y_2 , and the complement of the singularities in ∂Y_2 is a finite set of leaves. The set of singularities is also preserved by χ_0 . There is a length measurement on leaves in $\text{Interior}(Y_2)$ (induced by the unstable foliation being transverse to the stable), and there is $\lambda > 1$ such that χ_0 multiplies length by λ . Let x_i ($1 \leq i \leq r$) be the singularities. There is an integer $p_i \geq 3$ and a chart

$$\{z : |z| < 1\} \text{ if } x_i \in \text{Interior}(Y_2),$$

$$\{z : |z| < 1, \operatorname{Re}(z) \geq 0\} \text{ if } x_i \in \partial Y_2,$$

with x_i corresponding to 0 in the chart, such that the leaves of the foliation are locally the level sets of the multivalued function

$$\operatorname{Re}(z^{p_i/2}) \text{ if } x_i \in \operatorname{Interior}(Y_2),$$

$$\operatorname{Re}(z^{p_i-1}) \text{ if } x_i \in \partial Y_2.$$

Then by the Poincaré-Hopf Theorem,

$$\sum_{i=1}^r \left(\frac{p_i}{2} - 1 \right) = k - 1 \quad (1)$$

Since the number of singularities on each component of ∂Y_1 is the same (because χ_0 preserves the set of singularities) it follows from (1) that there must be exactly one singularity on each component of ∂Y_1 , and these are of the form $\chi_0^i x_1$ ($0 \leq i < k$) for some $x_1 \in \partial Y_1$ with $\chi_0^k x_1 = x_1$, and the p_i associated to x_1 must be 3, so that exactly one leaf ℓ in $\operatorname{Interior}(Y_2)$ ends at x_1 , and $\chi_0^k \ell = \ell$. Also, from (1) and the preservation of the foliation by χ_0 , we deduce that ℓ meets no other singularities. Also, since all infinite half-leaves of the unstable foliation in $\operatorname{Interior}(Y_2)$ are dense, $\tilde{\ell} = Y_2$. Since any measured foliation is recurrent, there is M_0 such that every segment of ℓ of length M_0 meets an element of Γ , where Γ is a fixed collection of disjoint simple closed curves on Y_2 cutting Y_2 into pairs of pants. It follows that the length of any segment ℓ_1 of ℓ is proportional to the diameter of any lift of it to $\tilde{Y}_2$.

3.4. Proof of the 1-sided Neighbourhood Theorem: finding Y_1 , T_1 , x_0 , χ_0 .

Since, by the Lefschetz Theorem, χ has a fixed point in Y , we can choose a connected subsurface Y_1 of Y which is invariant under χ up to isotopy, and such that no nontrivial nonboundary simple disjoint loop set in Y_1 is preserved by χ up to isotopy. Then we can assume $\chi Y_1 = Y_1$. Then χ cyclically permutes the components of ∂Y_1 , since χ cyclically permutes the components of $Y \setminus Y_1$. We can also assume there is a subgraph $T_1 \subset T$ with no free vertices such that $T_1 \subset Y_1$ and $T_1 \hookrightarrow Y_1$ is a homotopy equivalence. So there is $p : Y_1 \rightarrow T_1$ such that $p|_{T_1}$ is homotopic to the identity. Let $\tilde{Y}, \tilde{T}, \tilde{Y}_1, \tilde{T}_1$ be the universal covers of Y, T, Y_1, T_1 with $\tilde{T}_1 \subset \tilde{Y}_1 \subset \tilde{Y}$ and $\tilde{T}_1 \subset \tilde{T} \subset \tilde{Y}$. Let d be a fixed metric on $\tilde{Y}$ (lifted from a metric on Y) invariant under the action of the fundamental group $\pi_1(Y)$. Let $\tilde{h}, \tilde{\chi}, \tilde{p}$ be the lifts of h, χ, p so that a homotopy between $p \circ \chi$ and h (as maps from T_1 to T) lifts to a homotopy between $\tilde{h}$ and $\tilde{p} \circ \tilde{\chi}$. Then there is $M_1 > 0$ so that

$$d(\tilde{h}(x), \tilde{p} \circ \tilde{\chi}(x)) \leq M_1 \text{ for all } x \in \tilde{T}.$$

Since $\tilde{h}(\tilde{T}_1)$ is connected, for any $x_1, x_2 \in \tilde{T}_1$, $\tilde{h}(\tilde{T}_1)$ contains the unique arc between $\tilde{h}x_1$ and $\tilde{h}x_2$. Then we claim that

$$\tilde{T}_1 \subset \tilde{h}\tilde{T}_1.$$

For given $y \in \tilde{T}_1$, we can find $x_1, x_2 \in \tilde{T}_1$ such that $\tilde{p} \circ \tilde{\chi}(x_1), \tilde{p} \circ \tilde{\chi}(x_2)$ are separated by y and both distance $> M_1$ from y . Then y also separates $\tilde{h}x_1$ and $\tilde{h}x_2$. So $y \in \tilde{h}\tilde{T}_1$. Then a continuous map $\tilde{h}_0 : \tilde{T}_1 \rightarrow \tilde{T}_1$ is uniquely determined by the properties

$$\tilde{h}_0 = \tilde{h} \text{ on } \tilde{T}_1 \cap \tilde{h}^{-1}\tilde{T}_1,$$

$\tilde{h}_0$ maps each component of $\tilde{T}_1 \setminus \tilde{h}^{-1}\tilde{T}_1$ to a point.

Then $\tilde{h}_0$ descends to a map $h_0 : T_1 \rightarrow T_1$ which is homotopic to $\chi : T_1 \rightarrow Y_1$. So by the Lefschetz Fixed Point Theorem, h_0 has a fixed point x_0 . Then we can assume that $\chi(x_0) = x_0$, and that $\chi|_{T_1}$ and $h_0 : T_1 \rightarrow Y_1$ are homotopic via a homotopy fixing x_0 . Then we can find $\chi_0 : Y_1 \rightarrow Y_1$ as in the Pseudo-Anosov Proposition.

3.5. Proof of the 1-sided Neighbourhood Theorem: a reformulation.

It suffices to prove, for some lift H of h^k to $\tilde{T}$,

$$(3.5) \quad \tilde{T}_1 \subset \bigcup \{ \tau.W^u(x, H, U) : x \in \tilde{T}_1, Hx = x, U \text{ is a 1-sided neighbourhood of } x \text{ in } \tilde{T}_1, \text{ and } \tau \in \pi_1(T_1) \}.$$

For, projecting down, this proves the theorem.

3.6. Lemma.

It suffices to prove:

- (3.6) *for some closed arc I in $\tilde{T}_1$ projecting down to cover T_1 and some lift H' of h_0^k , $I \subset \bigcup \{ W^u(x, H', U) : x \in I, H'x = x \text{ and } U \text{ is a 1-sided neighbourhood of } x \text{ in } I \}$.*

Proof: Suppose (3.6) holds. Then let $\tau \in \pi_1(T_1)$ be such that $H' = \tau.\tilde{h}_0^k$, and let $H = \tau.\tilde{h}^k$. We claim that a), b) and c) hold. Clearly they suffice to prove (3.5), as is required.

- a) $W^u(x, H', U) \subset W^u(x, H, U)$ if $H'x = x$ and $\tilde{h}_0^i x = \tilde{h}^i x$ for $1 \leq i \leq k$,
- b) $W^u(x, H', U) = \{x\}$ if $H'x = x$ and $\tilde{h}_0^i x \neq \tilde{h}^i x$ for some $1 \leq i \leq k$,
- c) if, again, $\tilde{h}_0^i x \neq \tilde{h}^i x$ for some $1 \leq i \leq k$, then $\sigma.x \in W^u(x', H, U')$ for some $x' \neq \sigma.x$, $Hx' = x'$ and 1-sided neighbourhood U' of x' in I , and $\sigma \in \pi_1(T_1)$.

To see a), note that if V is any connected set in $\tilde{T}$, $\tilde{h}_0 V \subset \tilde{h}V$. We see b) as follows. If $x \in \tilde{T}_1$ and $H'x = x$ but $\tilde{h}^i x \neq \tilde{h}_0^i x$ for some $0 < i \leq k$, then we can find inductively $x_j \in \tilde{T}_1$ ($0 \leq j < k$) with $x_0 = x$, $\tilde{h}x_{j+1} = \tilde{h}_0x_{j+1} = x_j$.

Then for some $\zeta \in \pi_1(T_1)$, $\zeta.x_k \neq x$ but $H(\zeta.x_k) = H'(\zeta.x_k) = x$. Here, ζ is determined by τ , and by the automorphism of $\pi_1(T_1)$ which is itself determined by the homotopy class of $\tilde{h}$. Moreover, $W^u(x, H', U)$ is a point (for any 1-sided neighbourhood U of x), because, if i is the least integer > 0 with $\tilde{h}^i x \neq \tilde{h}_0^i x$, then i is also the least integer > 0 with $\tilde{h}^i x \notin \tilde{T}_1$, and there is a neighbourhood V of x with $\tilde{h}_0^i V = \tilde{h}_0^i x$, and then $H'V = x$. Now for c): by (3.6) there is $x' \in I$ and $\eta \in \pi_1(T_1)$ such that $\eta.x_k \in W^u(x', H', U')$ for some 1-sided neighbourhood of x' in I . Then $H(\eta.x_k) = H'(\eta.x_k) = \sigma.x \in W^u(x', H', U')$ for some $\sigma \in \pi_1(T_1)$. Now $\eta.x_k \neq \sigma.x$ because $\tilde{h}^i(\eta.x_k) \in \tilde{T}_1$ for $0 \leq i \leq k$, but this is not true for $\sigma.x$, because it is not true for x . So $H'(\eta.x_k) \neq \eta.x_k$ and $\eta.x_k \neq x'$. So $W^u(x', H', U')$ is not a point, so by b) $\tilde{h}_0^i x' = \tilde{h}^i x'$ for $0 \leq i \leq k$ and $W^u(x', H', U') \subset W^u(x', H, U')$ by a).

3.7.

There are two cases of the 1-sided Neighbourhood Theorem to prove, corresponding to cases 1 and 2 of the Pseudo-Anosov Proposition. In case 2, recall that χ_0 fixes a disc D round x_0 , and that $Y_2 = Y_1 \setminus D$. Then we choose a graph $T_2 \subset Y_2$ to be of the form $(T_1 \setminus D) \cup \partial D$, so that $T_2 \hookrightarrow Y_2$ is a homotopy inclusion, and we choose maps $h_2 : T_2 \rightarrow T_2$ and $p_2 : T_2 \rightarrow T_1$ so that $p_2 \circ h_2 = h_0 \circ p_2$, and p_2 is injective except for mapping ∂D to x_0 . Thus, p_2 extends to a map $Y_1 \rightarrow Y_1$ homotopic to the identity, and h_2 is homotopic to $g|_{T_2}$. Let $\tilde{p}_2$ be any lift of p_2 to $\tilde{Y}_1$. In case 2, it suffices to prove

- (3.7) for some closed arc I in $\tilde{T}_2$ projecting down to cover T_2 , and some lift H' of h_2^k , $I \subset \bigcup \{W^u(x, H', U) : x \in I, H'x = x \text{ and } U \text{ is a 1-sided neighbourhood of } x \text{ in } I\}$.

For suppose this is true. Then, by projecting by $\tilde{p}_2$ we find that (3.6) is true also for a suitable I and a suitable lift H' of h_0^k .

3.8. A further reduction.

We are now reduced to proving either (3.6) or (3.7), depending on whether we are in case 1 or 2. We claim that it suffices to prove

- (3.8) There is a closed arc I in $\tilde{T}_1$ (case 1) or $\tilde{T}_2$ (case 2) which projects to cover all of T_1 (case 1) or T_2 (case 2) and such that I has endpoints a, b , and there is a lift H' of h_0^k (case 1) or h_2^k (case 2) such that either $H'a = a$ or a separates $H'a$ and b , and either $H'b = b$ or b separates a and $H'b$, so that, in particular, $I \subset H'I$.

We shall complete the proof of the 1-sided Neighbourhood Theorem by proving that (3.8) holds in both cases 1 and 2. For the justification of this claim, we use the following lemma. Part (i) shows that I contains at least one fixed point. Then parts (ii) and (ii) are applied to subarcs of I between fixed points.

3.9. Lemma. *Let J be a closed arc in a tree with endpoints a, b , and let $H : T \rightarrow T$ be continuous. Then the following hold.*

(i) *If $H(a)$ is separated from b by a , and $H(b)$ is separated from a by b , then J contains a fixed point of H .*

(ii) *If $H(a) = a$ and $H(b)$ is separated from a by b , and J has no fixed interior point, then $J \subset W^u(a, H, J)$.*

(iii) *If $H(a) = a$, $H(b) = b$ and J has no fixed interior point, then $J \subset W^u(a, H, J) \cup W^u(b, H, J)$.*

Proof: (i) Inductively we can take $J = J_0$, and find intervals $J_{n+1} \subset J_n$ with $J_n \subset H(J_{n+1})$, and such that either J_n has a fixed endpoint for some n , or the conditions of (i) are satisfied by J_n for all n . In this latter case, $\cap J_n$ is either an interval with fixed endpoints, or is a fixed point.

(ii) By (i), for $a' \in \text{Interior}(J)$ sufficiently near a , a' is always separated from b by $H(a')$. Let c be the endpoint of $W^u(a, H, J) \cap J$ in $J \setminus \{a\}$. If $c \neq b$, then c separates $H(c)$ from b , since $H(c) \neq c$. By a), this cannot happen. So $c = b$. Then $b \in W^u(a, H, J)$, since $b \in H(\text{Interior}(J)) \subset W^u(a, H, J)$ if $c = b$. So $J \subset W^u(a, H, J)$, as required.

(iii) By (i), we can assume without loss of generality that a' is separated from b by $H(a')$ for $a' \in \text{Interior}(J)$ sufficiently near a . Let c be the endpoint of $W^u(a, H, J) \cap J$ in $J \setminus \{a\}$. If $c = b$, we are done. If not, then as in (ii), c separates $H(c)$ from b . Then we can complete the proof by applying (ii) with J replaced by the interval between c and b , and a replaced by b and b replaced by c .

3.10. Proof of (3.8) in Case 1.

By choice of $\tilde{h}$, we can ensure that $\tilde{h}_0$ fixes a preimage of x_0 in $\tilde{T}_1$. Take $H' = \tilde{h}_0^k$, and let the homotopy between χ_0^k and h_0^k lift to one between a map K and H' . Then K also fixes the same preimage of x_0 , so $K = \text{identity}$, since $\chi_0^k|Y_1 = \text{identity}$. So H' fixes all preimages of x_0 . We can take I to be any arc in $\tilde{T}_1$ with endpoints at preimages of x_0 and projecting to cover T_1 .

3.11. Proof of (3.8) in Case 2.

Let ℓ, x_1 be as in case 2 of the Pseudo-Anosov Proposition. Let $M > 0$, yet to be chosen. Let ℓ_1 be a segment of ℓ with endpoint at x_1 , such that

$$\text{length}(\chi_0^k(\ell_1) \setminus \ell_1) > M.$$

Let β be the component of ∂Y_1 containing x_1 . We can choose a lift $\tilde{\beta}$ of β in $\tilde{Y}_2$ and a lift K of χ_0^k so that all preimages of x_1 in $\tilde{\beta}$ are fixed by K . Let x'_1, x''_1 be two distinct preimages of x_1 in $\tilde{\beta}$, let ℓ'_1, ℓ''_1 be the lifts of ℓ_1 with endpoints at x'_1, x''_1 , let a', b' be the second endpoints of ℓ'_1, ℓ''_1 , and let J be

the arc in $\ell'_1 \cup \ell''_1 \cup \tilde{\beta}$ between a' and b' . Then $J \subset K(J)$, a' separates $K(a')$ from b' , b' separates $K(b')$ from a' , and by choice of M given $M_1 > 0$, we can ensure that

$$d(K(a'), a') > M_1, \quad d(K(b'), b') > M_1.$$

Since $T_2 \hookrightarrow Y_2$ is a homotopy equivalence, there is $q : Y_2 \rightarrow T_2$ which is homotopic to the identity as a map from Y_2 to Y_2 . Then the homotopy lifts to one between a map $\tilde{q}$ and the identity. Now we choose ℓ_1 so large that $q(J) = T_2$, which is possible by the Pseudo-Anosov Proposition. Then there is M_2 such that

$$d(Kx, H'\tilde{q}x) \leq M_2 \text{ for all } x \in \tilde{Y}_2,$$

where the homotopy between χ_0^k and h_0^k lifts to one between K and H' . We can choose $M_1 > M_2$. Then the arc I between $a = \tilde{q}a'$ and $b = \tilde{q}b'$ satisfies (3.8), as required.

Chapter 4. The Nonrational Lamination Map Theorem.

4.1.

The aim of this chapter is to prove the following theorem. For the definitions of invariant lamination, lamination map and so on, see Chapter 6 of [R2]. The definition of primitive can be found in Chapter 7 of [R2], and is quite complicated, but can mostly be forgotten.

The Nonrational Lamination Map Theorem.

Let ρ be a critically finite branched covering with a nondegenerate Levy cycle, which is also a lamination map, preserving a primitive invariant lamination L . Then there exist integers m, k , finite unions $A_0 \subset A_1 \subset \rho^{-1}A_0$ of leaf segments of L , and finite unions $B_0, B_1 \subset \rho^{-1}B_0$ of S^1 -segments, such that the following hold.

- a) Either $m = 1$, or mk is less than the minimum period of a periodic critical point of ρ .*
- b) Both $A_0 \cup B_0$ and $A_1 \cup B_1$ have m components, each homotopic in $\bar{C} \setminus X(\rho)$ to a k -holed sphere with boundary components nontrivial in $\bar{C} \setminus X(\rho)$.*
- c) Each component of A_0 contains a segment mapped into itself by ρ^{mk} , with orientation preserved.*
- d) For all $n \geq 0$, $\rho^{-n}B_0 \subset S^1$ and diameters of components tend to 0 as $n \rightarrow \infty$.*
- e) The sets $A_0 \cup B_0$ and $A_1 \cup B_1$ are homotopic in $\bar{C} \setminus X(\rho)$ via a homotopy which is constant on A_0 , and homotopes each component of $(A_1 \setminus A_0) \cup B_1$ into B_0 .*

4.2. Remark. If the theorem holds, we can inductively define A_n with $A_n \subset A_{n+1}$ and A_{n+1} a union of components of $\rho^{-1}A_n$, and A_n converges to a finite union A of leaves, each of which is fixed by ρ^{mk} , and $\bar{U}A$ is noncontractible in $\bar{C} \setminus X(\rho)$. It has m components, unless A contains compact geodesics.

4.3. Outline of Proof.

The theorem will be proved by choosing Y for $\rho = g$ as in the conclusion of the Fixed Holed-Sphere Theorem with certain additional properties, and then applying the 1-sided Neighbourhood Theorem to Y and some embedded graph T , where the isotopy class of the homeomorphism χ is given by a local inverse of ρ^m . Thus we have to choose $Y, T, h : T \rightarrow T$ with certain properties and deduce the theorem from this.

4.4. Recall of some basic definitions.

We shall need to refer to [R2] to some extent, particularly Chapters 6 and 7. Throughout the present chapter, L is a β -invariant lamination on $\bar{\mathbb{C}} \setminus K$, with lamination map ρ , which therefore preserves L and the Cantor set K , which is contained in S^1 . Recall that a *leaf segment* is the closure of a component of $\ell \setminus S^1$, for a leaf ℓ of L . A *polygon* is the closure P of a subset of $\bar{\mathbb{C}} \setminus S^1$ with $\partial P \subset (\cup L) \cup S^1$. A *gap* is a component of $\bar{\mathbb{C}} \setminus (K \cup (\cup L))$. A *gap polygon* is a polygon P such that $\text{Interior}(P)$ is a component of $G \setminus S^1$ for a gap G . A *side* of a gap G is a leaf ℓ with $\ell \cap P \neq \emptyset$ for some gap polygon P of G .

It was shown in [R2] - in particular, in 6.10 - that there are gap polygons Q_i ($1 \leq i \leq r$) and points a_i in K ($1 \leq i \leq q$) having the following properties, and so that ρ can be chosen to satisfy the following condition, given $\varepsilon' > 0$.

$$\rho^{-n}(S^1 \setminus ((\bigcup_{i=1}^q I_i) \cup (\bigcup_{i=1}^r \text{Interior}(Q_i))) \subset S^1 \text{ for all } n \geq 0,$$

where a_i is strictly preperiodic under ρ , and I_i is an interval in $S^1 \setminus K$ of length $< \varepsilon'$ and with endpoint at a_i , and $\rho(\cup Q_i) \subset \cup Q_i$. We shall need this because we shall need to consider families of leaf segments and S^1 -segments, which we shall want mapped to unions of leaf segments and to S^1 -segments by inverse images of ρ .

4.5. Essential Leaf Segments.

For suitable $\zeta > 0$, we say that a leaf segment ℓ is *essential* if some components of $\rho^{-n}\ell$ have diameter $> \zeta$ for infinitely many n . By the Length Lemma ([R2], 6.15), for suitable ζ , this is only possible if either

diameter(ℓ) $> \zeta$, or

diameter(ℓ) $\leq \zeta$ and ℓ has endpoints on opposite sides of, but close to, some a_i ($1 \leq i \leq q$).

From the definition (and for suitable ζ), if ℓ is not essential, then all components of $\rho^{-n}\ell$ are nonessential leaf segments, for all $n \geq 0$.

4.6. Properties of Y .

As we have said, we need to construct Y , T , h . We summarise the properties of Y we shall need, and which are proved in 4.7-4.10. That the first property is possible is simply a restatement of part of the Fixed Holed-Sphere Theorem.

1. Y is an open ℓ -holed sphere in $\bar{\mathbb{C}} \setminus X(\rho)$ for some $\ell \geq 2$, with each boundary component nontrivial, and for some local inverse S of ρ^m , Y and SY are isotopic in $\bar{\mathbb{C}} \setminus X(\rho)$, and S cyclically permutes boundary components up to isotopy in $\bar{\mathbb{C}} \setminus X(\rho)$.

2. Either (i) or (ii) holds, depending on whether the Expanding Fixed Set or Nonrational Adjacent case of the Fixed Holed-Sphere Theorem holds.

(i) $m = 1$ and $SY \subset Y$.

(ii) Here, m is such that $m\ell$ less than the minimum period of a periodic critical point of ρ , and $\partial Y = \partial_1 Y \cup \partial_2 Y$, where $S\bar{Y} \cap \partial_1 Y = \phi$, and $S\partial_2 Y \cap \bar{Y} = \phi$. Each component C of $SY \setminus Y$ intersects only one component of each of $\partial_2 Y$, $S\partial_2 Y$.

For (i) see 4.7, and (ii) is a restatement of the Fixed Holed-Sphere Theorem.

3. Either (i) or (ii) holds, depending, again, on the case of the Fixed Holed-Sphere Theorem.

(i) ∂Y is a finite union of leaf segments and S^1 -segments.

(ii) $\partial_1 Y$ is a finite union of leaf segments and S^1 -segments, $\partial_2 Y$ is a finite union of S^1 -segments.

For (i), see 4.7. For (ii), see 4.9.

Note. We shall write $S^n Y$ even when $SY \setminus Y \neq \phi$, meaning the local inverse defined on $S^{n-1}Y$ by lifting the inductively defined homotopy between $S^{n-2}Y$ and $S^{n-1}Y$.

4. $S^i Y \cap \{a_1 \dots a_q\} = \phi$ for all $i \geq 0$, and if b is a component of $\partial Y \cap S^1$, then $b \cap \bigcup_{i=1}^r Q_i = \phi$ so that $S^n b \subset S^1$ for all n .

See 4.7.

5.

$$\bar{Y} = \bigcup_{i=1}^v Y_i,$$

where the sets $\text{Interior}(Y_i)$ are disjoint, and, for some $0 \leq t \leq u \leq v$, the following hold.

a) $\partial Y_i \setminus \partial Y \subset S^1$, and if b is a component of $\partial Y_i \cap S^1$, $b \cap \bigcup_{i=1}^r Q_i = \phi$, so that $S^n b \subset S^1$ for all n .

b) For $i > t$, Y_i is a single polygon not containing any periodic gap polygon such that ∂Y_i contains exactly two essential leaf segments for $t < i \leq u$, and none for $u < i \leq v$.

c) For $i \leq t$, Y_i is a finite connected union of polygons, and there is a finite connected union $Y'_i \subset Y_i$ of gap polygons such that each maximal polygon of Y_i contains exactly one polygon of Y'_i , $(Y_i \setminus Y'_i) \cap S^1 \cap \bigcup_{i=1}^r Q_i = \phi$, and there is a bijection $\tau : \{1 \dots t\} \rightarrow \{1 \dots t\}$ with

$$Y'_{\tau(i)} \subset SY'_i.$$

d) $SY_j \cap Y_i = \phi$ for $i, j \leq t$ and $i \neq \tau(j)$, and $SY_j \cap Y_{\tau(j)}$ is connected.

See 4.10 (and 4.7-4.9).

4.7. Lemma.

Y can be chosen so that either 1 or 2 holds, where S is a local inverse of ρ^m .

1. $m = 1$, ∂Y is a finite union of leaf segments and S^1 -segments, and $SY \subset Y$.
2. $\partial Y = \partial_1 Y \cup \partial_2 Y$, where $\partial_i Y$ is a finite union of leaf segments and S^1 -segments,

$$S\bar{Y} \cap \partial_1 Y = \phi, S\partial_2 Y \cap \bar{Y} = \phi,$$

and each component of $SY \setminus Y$ has one component of intersection with each of $\partial_2 Y$, $S\partial_2 Y$.

Furthermore, we can ensure that $\partial Y \cap S^1 \cap Q_i = \phi$ ($1 \leq i \leq r$) and $S^n Y \cap \{a_1 \dots a_q\} = \phi$ for all $n \geq 0$.

Proof:

Suppose the Expanding Fixed Set case of the Fixed Holed-Sphere Theorem holds. Then the variation of 2.3 implies Y can be chosen to contain all leaves and finite-sided gaps of L it intersects, and so that $SY \subset Y$ and $S^p \bar{Y} \subset Y$ for some p . For any finite connected union of leaf segments, or arc in a finite-sided gap has the property of γ in the statement of the variation. Then we can approximate $S^i Y$ ($0 \leq i < p$) by a set Y_i whose boundary is a finite union of leaf segments, components of intersection with gap polygons in infinite-sided gaps and their sides, and S^1 -segments, such that $\bar{Y}_{i+1} \subset Y_i$ for $i \leq p-2$, $S\bar{Y}_{p-1} \subset Y_0$ and $\partial Y_i \cap \partial Q_j \cap S^1 = \phi$. Then $Y' = \bigcap_{i=0}^{p-1} Y_i$ satisfies $S\bar{Y}' \subset Y'$. Let Y be obtained from Y' by removing all components of intersection with gap polygons in infinite-sided gaps and their sides which are not complete gap polygons. Then $SY \subset Y$. We can ensure $S^n Y \cap \{a_1 \dots a_q\} = \phi$ for all $n \geq 0$ by replacing Y by $S^n Y$ for a suitable $n \geq 0$, because the a_i are strictly preperiodic.

Suppose the Nonrational Adjacent case of the Fixed Holed-Sphere Theorem holds. Then the variations of 2.3 ensure that we can choose Y so that $\partial_1 Y$, $\partial_2 Y$ consist of finitely many leaf and S^1 -segments, and do not intersect $S^1 \cap Q_i$ for any i . Then $\partial_1 Y$, $\partial_2 Y$ have all the required properties, provided, as in case 1, we replace Y by $S^n Y$ if necessary.

4.8. Lemma.

Y can be chosen so that the periodic gap polygons contained in $S^n Y$ are the same for all $n \geq 0$, and such that no two periodic gap polygons P , Q in Y lie in the same component of $Y \setminus S^1$.

Proof: Since the lamination L is primitive, the total number of periodic gap polygons is finite. (See 6.15, 7.2 and 7.3 of [R2]. A periodic thin gap polygon lies in a periodic gap bounded by periodic segments on some of the finitely many periodic sides. But in a primitive lamination, there is a bound on the number of periodic segments on any leaf.) If $SY \subset Y$, the number of periodic gap polygons

in $S^n Y$ obviously decreases with n , so replacing Y by a suitable $S^n Y$, we can assume the number is always constant. If Y satisfies the Nonrational Adjacent case of the Fixed Holed-Sphere Theorem, so that $SY \setminus Y \neq \phi$, let p be the maximal period of a periodic gap polygon. If some periodic gap polygon is in $SY \setminus Y$, then for some $1 \leq n \leq p$, $S^n(SY \setminus Y) \cap Y \neq \phi$ - but, by the variations of 2.3 of the Fixed Holed-Sphere Theorem, Y can be chosen so that this does not happen.

Finally, suppose for contradiction that P, Q are in the same component of $S^n Y \setminus S^1$ for all n . Then P, Q are finite-sided, since $S^n \mid Y$ is a homeomorphism for all n . We can assume there are no periodic gap polygons between P and Q in the same component of $Y \setminus S^1$. But the region between P and Q must be mapped back into itself by some S^n , so it must be a single periodic leaf segment, which is impossible, because gaps are never immediately adjacent in a primitive lamination. So, again, replacing Y by $S^n Y$ if necessary, Y has the required properties.

4.9. Lemma.

In the case when $\partial Y = \partial_1 Y \cup \partial_2 Y$, Y can be chosen so that $\partial_2 Y \subset S^1$.

Proof: Let C be a component of $SY \setminus \bar{Y}$. Then by the variations of 2.3, the distance between $\bar{C} \cap \partial_2 Y$ and $\bar{C} \cap S\partial_2 Y$ can be assumed so large that no maximal polygon in C , or union of polygons intersecting some Q_i , has sides in both $\partial_2 Y$ and $S\partial_2 Y$. Then we can find an S^1 -arc in C , which intersects no Q_i , separating $\bar{C} \cap \partial_2 Y$ and $\bar{C} \cap S\partial_2 Y$. Then we replace Y by SY cut off by S^1 -arcs in all such components C .

4.10.

4.7 to 4.9 show that Y can be chosen satisfying 1 to 4 of 4.6, and part of 5 of 4.6: we can write $\bar{Y} = \cup_{i=1}^r Y_i$ so that Y_i is a single polygon not containing a periodic gap polygon for $i > t$, and Y_i has the properties listed in 5 of 4.6 for $i \leq t$, except for 5b) and 5d). Now we claim that, if we replace Y by $S^n Y$ for a suitable $n \geq 0$, and rename and renumber the Y_i , but keeping the name Y_i ($i \leq t$) for the union of polygons of $\bar{Y}$ intersecting Y'_i , then 5 of 4.6 is satisfied in its entirety. For gap polygons are dense and all eventually periodic, so we can ensure that, for $i > t$, Y_i does not contain any specified gap polygon, by choosing n large enough. So then, given $\zeta > 0$, we can ensure that ∂Y_i has at most two leaf segments of length $> \zeta/2$, and if at most one has length $> \zeta/2$, they all have length $< \zeta$. Thus we can ensure that ∂Y_i contains either exactly two essential leaf segments, or none, for $i > t$, and, renumbering, 5a)-c) of 4.6 are satisfied. we can also assume that $\cup_{i=1}^t (Y_i \setminus Y'_i) \cap \cup_{i=1}^t SY'_i = \phi$, so that 5d) is also satisfied, as required.

4.11. Properties of U , U' .

Let $U \subset \bar{U} \subset U'$ be open sets in $\bar{Y}$ satisfying the following - which are possible.

$$a) \quad \bigcup_{i>u} Y_i \subset U,$$

and, in the Nonrational Adjacent case,

$$\partial_2 Y \subset U.$$

b) For $t < i < u$, $Y_i \cap U$ has two components, and $\partial(Y_i \setminus U) \setminus \partial U$ has two components, which contain $\bar{Y}_i \cap S^1$, and the essential leaf segments in ∂Y_i run between these two components.

c) For $i \leq t$, each component of $Y_i \cap U$ contains exactly one component of $\partial Y_i \cap S^1$.

d) Similar properties to a), b) and c) hold for U' .

$$e) \quad S\bar{U} \cap Y \subset U'.$$

f) Each component of $SY \setminus Y$ intersects a unique component of U' .

4.12. Properties of T .

We can define a graph $T \subset Y$ satisfying the following.

a) $T \hookrightarrow Y$ is a homotopy equivalence.

b) T has no vertex in $Y_i \setminus U$ for $t < i \leq u$, but has one edge crossing between the components of U bounding Y_i .

c) For $i \leq t$, $T \cap Y_i \subset Y'_i$, T has exactly one vertex v_i in $Y'_i \setminus U$ with $Sv_i \in Y'_{\tau(i)} \setminus U'$, and exactly one edge of T runs from v_i into each bounding component of U .

d) No arc of $ST \setminus U'$ has both endpoints in the same component of U .

4.13. Definition of π , h .

We can define $\pi : ST \rightarrow T$ satisfying the following.

a) π maps each component of $ST \cap U \setminus \bigcup_{i=1}^t Y_i$ into the component of U' it is contained in, and each component of $ST \setminus Y$ into the component of U' it intersects.

b) π maps $ST \setminus (U' \cup \bigcup_{i=1}^t Y_i)$ to $T \setminus U$, and is monotone restricted to each arc in $ST \setminus (U \cup \bigcup_{i=1}^t Y_i)$.

$$c) \quad \pi Sv_i = v_{\tau(i)}.$$

d) Any component of $ST \cap (\bigcup_{i=1}^t Y_i \setminus U)$ whose closure intersects only one component of U' is mapped by π into that component of U' . For $i \leq t$, π is monotone restricted to each arc in $ST \cap Y_i \setminus U'$ which has endpoints in different components of U' , and maps it to an arc in $Y_i \cap T$ with endpoints in the same components of U .

Then we define

$$h = \pi \circ S.$$

4.14. Properties of h .

From the definition of h , we obtain the following.

a)
$$h(U) \subset U.$$

b) If $t < j \leq u$ and $i \leq u$, then $h(T \cap Y_i \setminus U)$ contains $T \cap Y_j \setminus U$ if it intersects it. This happens if and only if $SY_i \cap Y_j$ separates the essential leaf segments in ∂Y_j .

c) For $i \leq t$, $h(v_i) = v_{\tau(i)}$, and if $j \leq t$ also,

$$h(T \cap Y_i \setminus U) \cap Y_j = \phi,$$

unless $j = \tau(i)$, when $h^{-1}(Y_j \cap T \setminus U) \cap T \cap Y_i \setminus U$ is connected, and then, if b is a component of $Y_i \setminus U \setminus \{v_i\}$, $h(b)$ contains exactly one component of $Y_j \setminus U \setminus \{v_j\}$ and intersects no other.

Now the 1-sided Neighbourhood Theorem will imply the existence of k with $2 \leq k \leq \ell$ and a subgraph T_1 of T with

d)
$$T_1 \setminus U \subset \bigcup \{W^u(x, h^k, V) : x \in T_1 \setminus U, h^i x \in T_1 \text{ for all } i, h^k x = x,$$

V is a 1-sided neighbourhood of x in $T_1\}$.

Then the properties of π imply that

e) if $h^k x = x$ and $W^u(x, h^k, V) \neq \phi$ for some $x \in T_1 \cap Y_i \setminus U$, some $i \leq t$ and for some V , then $b \subset S^k b$ for each component b of $\partial Y_i' \cap (\cup L)$. Moreover, given $i \leq t$, there must be such an x, V , unless every arc of $T_1 \cap Y_i \setminus U'$ is in $h(T_1 \setminus Y_i)$, that is, unless, for any maximal arc b of $T_1 \cap Y_i$, there is an arc of $S(T_1 \cap Y \setminus Y_i) \cap Y_i$ running parallel to it throughout its length.

4.15. Proof of the Nonrational Lamination Map Theorem.

We shall find A_0 such that exactly one connected leaf piece from A_0 runs parallel to each arc in $T_1 \cap (\bigcup_{i=1}^u Y_i \setminus U)$. Then we shall join the endpoints of the leaf pieces of A_0 by S^1 -arcs whose union we denote by B_0 , so that $A_0 \cup B_0$ is homotopic to T_1 in $\bar{\mathcal{C}} \setminus X(\rho)$. We shall have $\rho^{-n} B_0 \subset S^1$ for all n , since each component of B_0 is a union of a subset of $\cup Y_i \cap S^1$ and of intervals in S^1 between points of $Y_j \cap S^1$ ($j > u$). Thus, $\rho^{-n} B_0 \cap (\cup Q_i \cup \{a_1..a_q\}) = \phi$ for all

$n \geq 0$. The homotopy between $A_0 \cup B_0$ and $A_1 \cup B_1$ is obtained by constructing a homotopy between $A_0 \cup B_0$ and T_1 which preserves U , $Y \setminus U$, constructing a similar homotopy between ST_1 and $A_1 \cup B_1$, and constructing yet another similar homotopy between π and the identity. So it remains to construct A_0 . From properties b), d), e) of h in 4.14, the proof of the theorem will be completed after the following lemma.

4.16. Lemma.

If $t < i \leq u$ and $x \in Y_i \cap T_1 \setminus U$, $h^k x = x$, $h^j x \in T_1$ for $0 \leq j \leq k$, and $W^u(x, h^k, V) \neq \emptyset$, then there is a leaf segment ℓ in Y_i separating the essential leaf segments in ∂Y_i such that $\ell \subset S^k \ell$, and $S^k \mid \ell$ preserves orientation.

Proof:

Let $x \in Y_i \cap T_1 \setminus U$, $t < i \leq u$, $h^k x = x$, $h^j x \in Y_{i_j} \cap T_1 \setminus U$ with $i_0 = i_k = i$. Then $t < i_j \leq u$ for all j . Inductively, let $Q_0 = Y_i$, and let Q_{j+1} be the component of $SQ_j \cap Y_{i_{j+1}}$ in the component of $SY_{i_j} \cap Y_{i_{j+1}}$ which contains $Sh^j x$ for $0 \leq j < k$. Then, for $n \geq k$, let Q_{n+1} be the unique component of $SQ_n \cap Q_{n+1-k}$. Then inductively, using property b) of h in 4.13, we see that Q_{nk+j} separates the essential leaf segments in ∂Y_{i_j} . Then $\cap Q_{nk}$ must be a leaf segment ℓ with $\ell \subset S^k \ell$, because otherwise it is a periodic gap polygon in Y_i , and there are none. If $W^u(x, h^k, V) \neq \emptyset$ for some V , then $S^k \mid \ell$ preserves orientation. This is because, for all $j \leq k$, the images under S and h of the endpoints of the arc of $T \cap Y_{i_j} \setminus U$ containing $h^j x$ lie in the same components of U , and the endpoints of the arc of $S^{j+1} \ell$ in Q_{j+1} also lie in these same components of U .

4.17. End of proof of the Theorem.

It suffices to choose $A_0 \subset Y$ so that a leaf segment from A_0 intersects $Y_i \setminus U$ for all $t < i \leq u$, and some connected union of leaf segments in A_0 runs parallel to each component of $\partial Y'_i$, for all $i \leq t$. Now fix $i \leq u$, and suppose $W^u(x, h^k, V) \neq \emptyset$ for some $x \in T_1 \cap Y_i \setminus U$, and V . If $i \leq t$ let $\partial Y'_i \cap (\cup L) \subset A_0$, and if $t < i \leq u$, let A_0 include a leaf segment ℓ as in the lemma, and let A_0 also contain a finite connected union of segments containing ℓ which is contained in Y , has endpoints in U , and runs parallel to as many arcs of $T_1 \setminus U$ as possible. Then A_0 has the required properties.

Chapter 5. Proofs of the Admissible Boundary Theorem and the Admissibility Proposition.

5.1.

The theorems to be proved in this chapter are stated in Chapter 1 of [R2], and will be restated here, but it will not be possible to explain all the terms involved, for which the reader should refer to Chapters 6 and 7 of [R2], in particular, for a full account of invariant and parameter laminations. The main result of the last chapter, the Nonrational Lamination Map Theorem, was essentially a result about invariant laminations (or their lamination maps). The task, now, is to transfer this to a result about parameter laminations.

5.2. Invariant sets of complete geodesics.

We recall, at least, that an invariant lamination L is a geodesic lamination on $\bar{\mathbf{C}} \setminus K$, where K is a Cantor subset of S^1 , where $K = K_r$ depends on a parameter r (which is an odd denominator rational). Also, there is a fixed lamination called (somewhat incongruously at this stage) $\bar{L}_r$ such that $L \cup \bar{L}_r$ is always a lamination, and L contains all but countably many leaves of $\bar{L}_r$. The sets S^1 , K and $\bar{L}_r$ are invariant under a critically finite degree two branched covering $\bar{s} = \bar{s}_r$ which fixes ∞ . For precise definitions of K , $\bar{L}_r$, $\bar{s}$, see Chapter 1 of [R2].

The invariance is actually β -invariance for some path $\beta : [0, 1] \rightarrow \bar{\mathbf{C}} \setminus (K \cup (\cup \bar{L}_r))$, and uses a homeomorphism σ_β defined in terms of β . (See [R2] 1.7.) We recall that, in [R2] 6.5 we put extra conditions on β in order to ensure that, for a complete geodesic ℓ in a β -invariant lamination, $(\sigma_\beta \circ \bar{s})^{-1}\ell$ could be easily modified to a set of geodesics. We also need to do something similar here for finite geodesics, that is, geodesics with finitely many S^1 -crossings, which may be incomplete. This means making modifications to β which are similar to those of 6.5 of [R2], but slightly less technical. We describe these now.

5.3. The Minimal Intersection Property.

Let $\beta : [0, 1] \rightarrow \bar{\mathbf{C}} \setminus (K \cup (\cup \bar{L}_r))$ with $\beta(0) = \infty$ and let A be a union of finite disjoint geodesic arcs in $\bar{\mathbf{C}} \setminus (K \cup (\cup \bar{L}_r))$ with any endpoints in S^1 . Then after isotopy in $\bar{\mathbf{C}} \setminus (K \cup (\cup \bar{L}_r))$ keeping $\beta(0) = \infty$ and fixing $\beta(1)$, β can be assumed to have the following properties.

- a) If c, d are adjacent points of $\beta^{-1}(S^1)$, then $\beta(c), \beta(d)$ lie in different components of $S^1 \setminus K$, and β is transverse to S^1 .
- b) No sub-path of β can be homotoped into A by a homotopy in $\bar{\mathbf{C}} \setminus K$ fixing the sub-path endpoints, β is transverse to A , and the number of points of

$(0, 1) \cap \beta^{-1}(A)$ cannot be reduced by homotopies fixing $\beta(1)$ and $\beta(0) = \infty$, while keeping property a).

Let β_u be any homotopy through paths in $\bar{\mathbf{C}} \setminus (K \cup (\cup \bar{L}_r))$ satisfying a) and b) such that $\beta_0 = \beta$, $\beta_u(0) = \beta(0) = \infty$ and $\beta_u(1) = \beta(1)$ for all u , and such that $\beta_u(t) \in A$ only if $\beta(t) \in A$. Now we can state condition c).

c) If $\beta(t) \in A$, then $\beta(t) \notin S^1$, $\beta_u(t) \in A$ and

$$\#\{v < t : \beta(v) \in S^1\} \leq \#\{v < t : \beta_u(v) \in S^1\}$$

for all u .

The conditions b), c) subject to a) ensure that $(\sigma_\beta \circ \bar{s})^{-1}A$ has the least possible number of intersections with S^1 , up to homotopy fixing the endpoints of $(\sigma_\beta \circ \bar{s})^{-1}A$ and preserving K and $\bar{L}_r$.

Then (β, A) has the *minimal intersection property* if conditions a) to c) hold. A *good isotopy* will be one which preserves S^1 , K and $\bar{L}_r$.

Note the following.

1. The image of (A, β) under any good isotopy also has the minimal intersection property.
2. If A can be isotoped into A' by a good isotopy, then there is β' such that (A', β') has the minimal intersection property, and such that (A, β) can be isotoped into (A', β') by a good isotopy.
3. For any finite set A of finite geodesics in a β -invariant lamination with β satisfying the conditions of 6.5 of [R2], (A, β) has the minimal intersection property.

5.4. First Invariant-implies-Parameter Theorem. *In what follows, all inclusions and equalities are up to good isotopy. Let (A_0, β) have the minimal intersection property, with $\beta(1)$ in a component μ_0 of A_0 , and suppose that*

$$A_0 \subset (\sigma_{\beta'} \circ \bar{s})^{-1}A_0,$$

where β' is an arbitrarily small extension (or restriction) of β . If $x \notin K$ is an endpoint of A_0 , let $\tau(x)$ be the endpoint (if it exists) of A_0 such that $(\sigma_{\beta'} \circ \bar{s})^{-1}(\tau(x))$ is isotoped to x by the good isotopy. Suppose also that, for any endpoint $x \notin K$, $\tau^n(x)$ is not defined for all n .

Then there are γ , and a γ -invariant lamination L , such that $A_0 \subset L$, μ_0 is contained in the minor leaf or a side of the minor gap of L , and (A_0, β') or $(A_0, \beta) = (A_0, \gamma)$ up to good isotopy, depending on whether or not L has a minor gap.

The minor leaf of L is eventually periodic, and strictly preperiodic if L does not have a minor gap. Moreover, no half leaf of L has infinitely many successive intersections with periodic components of $S^1 \setminus K$.

Proof: Again, all inclusions and equalities are up to good isotopy. Write $\beta_0 = \beta'$. Inductively, define β_n, A_n for $n \geq 1$ with the following properties.

1. $A_{n+1} = (\sigma_{\beta_n} \circ \bar{s})^{-1} A_n$
2. $(A_n, \beta_n) \subset (A_{n+1}, \beta_{n+1})$
3. (A_n, β_n) has the minimal intersection property.

We can clearly define A_1, β_1 . Suppose A_k, β_k have been defined with the correct properties for $k \leq n$ and $n \geq 1$, and define A_{n+1} by 1. Then since $(A_{n-1}, \beta_{n-1}) \subset (A_n, \beta_n)$, we have $(\sigma_{\beta_{n-1}} \circ \bar{s})^{-1} A_{n-1} \subset (\sigma_{\beta_n} \circ \bar{s})^{-1} A_n$, and hence $A_n \subset (\sigma_{\beta_n} \circ \bar{s})^{-1} A_n$. Then by 5.3.1 and 2 we can find a β_{n+1} so that $(A_n, \beta_n) \subset (A_{n+1}, \beta_{n+1})$.

Then the condition on $\{\tau^n(x)\}$ implies that, if x is an endpoint of some A_n , then either $x \in K$ or x is interior to A_m for some $m > n$. It follows that A_n converges to a lamination L , containing A_0 . We can also assume that β_n converges to γ' . Here, γ' might contain points of K which are endpoints of intervals of $S^1 \setminus K$, but can be perturbed to γ in $\bar{\mathcal{C}} \setminus (K \cup (\cup \bar{L}))$ such that L is γ -invariant. Note that A_0 contains all the periodic segments in the full orbit of A_0 , and L is the closure of this full orbit. Hence no half leaf in L has an infinite succession of periodic segments, as required.

Let ℓ_0 be the component of A_0 containing $\beta(1)$ and let ℓ_{i+1} be defined inductively as a component of A_{i+1} which is a component of $(\sigma_{\beta_i} \circ \bar{s})^{-1} \ell_i$. Obviously, ℓ_0 in L is always in an eventually periodic leaf, and in a periodic leaf if and only if $\ell_0 \subset \ell_n$ with orientation preserved for some n , for some choice of $\{\ell_i\}$. If this happens, no other component ℓ'_n of A_n can run parallel to ℓ_n for the whole length of ℓ_n on the same side as $\beta_n(1)$. For if so, let ℓ'_i ($0 \leq i \leq n$) be defined similarly to ℓ_i , so that ℓ'_{i+1} is a component of $(\sigma_{\beta_i} \circ \bar{s})^{-1} \ell'_i$. Suppose ℓ_{i+1} and ℓ'_{i+1} run parallel along their entire lengths. Then $\sigma_{\beta_i}^{-1} \ell_i$ and $\sigma_{\beta_i}^{-1} \ell'_i$ run parallel, and so do ℓ_i and ℓ'_i . Then, by induction, $\sigma_{\beta'}^{-1} \ell_0$ and $\sigma_{\beta'}^{-1} \ell'_0$ run parallel along their entire lengths. This is impossible, because $\beta'(1)$ separates ℓ_0 and ℓ'_0 . It follows that β_k can be chosen so that (A_k, β_k) converges to (L, γ') , modifying to (L, γ) , with $\gamma(1)$ in a minor gap of L . Thus, L has all the required properties.

5.5.

We recall that, in [R2], $\tilde{U}$ denoted the universal cover of the union of component of $\bar{\mathcal{C}} \setminus (K \cup (\cup \bar{L}_r))$ containing ∞ and some boundary leaves, with $0 \in \tilde{U}$ projecting to ∞ . If $\beta : [0, 1] \rightarrow \bar{\mathcal{C}} \setminus (K \cup (\cup \bar{L}_r))$ is a path with $\beta(0) = \infty$, then $\tilde{\beta}$ denotes the lift to $\tilde{U}$ of β with $\tilde{\beta}(0) = 0$. Given a β -invariant lamination L with minor gap, the lift to $\tilde{U}$ containing $\tilde{\beta}(1)$ is also called the minor gap, and similarly for minor leaves: the minor leaf in $\tilde{U}$ either contains $\tilde{\beta}(1)$, or is intersected by $\tilde{\beta}$ and in the boundary of the minor gap, if this exists.

We recall the Parameter Laminations Theorem (1.16 of [R2]) which says that minor leaves and boundaries of minor gaps of primitive invariant laminations never intersect transversally. Primitivity is no real restriction. As in the statement and proof of the First Invariant-implies-Parameter Theorem, inclusions and equalities for sets of geodesic pieces are up to good isotopy, in the following theorem.

Second Invariant-implies-Parameter Theorem.

Let (A_0, β) and β' be as in the First Invariant-implies-Parameter Theorem. Let L' be a primitive γ' -invariant lamination and let $\tilde{\gamma}'$ be the lift of γ' with $\tilde{\gamma}'(0) = 0$. Suppose that there is an isotopy preserving K , $\bar{L}_r$ and A_0 of β' to γ' . Suppose the minor gap of L' - if it exists - is not both periodic and nonsimply-connected. Then any transverse intersections between A_0 and the minor leaf or sides of the minor gap of L' can be removed by good isotopy, and one of the following holds.

1. Each component of A_0 is contained in L' or a finite-sided gap of L' .
2. There is a γ -invariant lamination L containing A_0 and with all the other properties of the First Invariant-implies-Parameter Theorem, such that $L' \subset L$. We can choose L so that either of the following holds. The minor leaves of L , L' are the same, or at least one component of A_0 is contained in the forward orbit of the minor leaf or sides of the minor gap of L .

Proof: The proof is similar to that of the First Invariant-implies-Parameter Theorem, but we start by fixing a finite set C_0 of finite geodesics in the forward orbit of sides of the minor gap of L' - if this exists - or in the minor leaf of L' , such that (C_0, γ') has the minimal intersection property,

$$C_0 \subset (\sigma_{\gamma'} \circ \bar{s})^{-1} C_0,$$

and the minor leaf and sides of the minor gap (if the minor gap exists) are in the full orbit of C_0 . Now we construct inductively $A_n \subset A_{n+1}$, $C_n \subset C_{n+1}$, with

$$A_{n+1} = (\sigma_{\beta_n} \circ \bar{s})^{-1} A_n, \quad C_{n+1} = (\sigma_{\beta_n} \circ \bar{s})^{-1} C_n,$$

where the definition of β_n is slightly different from that in the First Theorem in some cases, and we do not define C_m for $m \geq n$ if C_n intersects A_n transversally. For $n \geq 0$, β_n is always defined so that either (A_n, β_n) or $(A_n \cup C_n, \beta_n)$ has the minimal intersection property, depending on whether or not C_i intersects A_i transversally for some $i \leq n$. Similarly, β_n and β_{n+1} have the same intersections with A_n or $A_n \cup C_n$. Also, $\beta_n(1)$ is always in the minor gap of L' . The lamination L is then obtained by taking the limit of A_n or $A_n \cup C_n$, depending on whether or not C_n intersects A_n transversally. The Parameter Laminations Theorem (1.16 [R2]) and (7.1, 7.3 [R2]) then show that, in fact, C_n does not intersect A_n transversally unless the minor gap of L' is periodic nonsimply-connected. Every leaf of L is contained in L' or a finite-sided gap of L unless the minor gap of L' is periodic (and hence infinite-sided).

There is a certain amount of choice in the construction. If some A_m intersects C_m transversally or the minor gap of L' , then for some N , β_n must always be close to the same component of $C_N \subset C_n$ or $A_N \subset A_n$ for $n \geq N$. Otherwise, β_n must be close to the same component of $C_N \subset C_n$ for $n \geq N$ for some N . We can choose the component of A_N to contain a component of A_0 , if the minor gap of L' is periodic. We can satisfy these conditions, so we are done.

5.6. Piecewise-geodesic Fixed Holed-spheres.

Recall that $\bar{s}$ is a critically finite branched covering, and $X(\bar{s}) \setminus \{\infty\} = X_1(\rho) = \{\rho^i(0) : i \geq 0\}$ for any branched covering lamination map ρ (for an invariant lamination containing $\bar{L}_r$).

Given an invariant lamination L with lamination map ρ , let A_0, B_0 be the sets proved to exist in the Nonrational Lamination Map Theorem when L is primitive and ρ has a nondegenerate Levy cycle. It is clear, by considering the boundary of the set $A_0 \cup B_0$, that the converse is true: if ρ is a critically finite branched covering, and such that A_0, B_0 exist, then ρ has a nondegenerate Levy cycle. The important thing to remember is that, if m is the period of 0 under $\bar{s}$ (and ρ) then all segments of A_0 are eventually periodic of eventual oriented period $\leq m$, and if $A_0 \cup B_0$ is not connected, the eventual oriented period is strictly less than the period of any periodic critical point. (I apologise for the change in use of m . We now revert to the letter conventions established in Chapter 1 of [R2]. The lettering in Chapters 1 to 4 of this paper is internally consistent as far as I can manage it, and that is the best I can do.) We shall call $A_0 \cup B_0$ a *piecewise-geodesic fixed holed-sphere*, if it is connected, and, in general, a *piecewise-geodesic periodic holed-sphere*. Recall that $A_n \cup B_n$ is the component of $\rho^{-n}(A_0 \cup B_0)$ containing A_0 , and the properties satisfied by $A_0 \cup B_0$ imply that $\bar{U}A$ is noncontractible in $\bar{\mathbf{C}} \setminus X_1(\rho) = \bar{\mathbf{C}} \setminus (X(\bar{s}) \setminus \{\infty\})$.

We also have the following, using the same notations as in the First Invariant-implies-Parameter Theorem and its proof.

Lemma. *Let (A_0, β) and L be as in the First Invariant-implies-Parameter Theorem. Let ρ be the lamination map of L . Also, let B_0 be a finite union of intervals in $S^1 \setminus K$ with endpoints in A_0 , such that the forward orbits of the components of $S^1 \setminus K$ intersected by β are not intersected by B_0 . Then $\rho^{-n}(B_0) \subset S^1$ for all $n \geq 0$.*

Similarly, let $(A_0, \beta), L'$, with lamination map ρ' , be as in the Second Invariant-implies-Parameter Theorem, with the same hypotheses as above. If the first conclusion of that theorem holds, $\rho'^{-n}(B_0) \subset S^1$ for all n . If the second conclusion holds, L can be chosen so that, in addition to the other properties, $\rho'^{-n}(B_0) \subset S^1$ for all n .

Proof: In both the First and Second Theorems, (A_n, β_n) has the minimal intersection property. It suffices that, for all n , $\beta_n \cap B_n = \emptyset$, where $B_{n+1} = (\sigma_{\beta_n} \circ \bar{s})^{-1} B_n$. In the case of the First Theorem, this is obvious, by the condition on β , since $B_n = (\sigma_{\beta_n} \circ \bar{s})^{-1} B_{n-1} = (\sigma_\beta \circ \bar{s})^{-n} B_0 = \bar{s}^{-n} B_0$. In the case of the Second Theorem, it is also clear, unless some component of A_n is in the minor gap of L' . In that case, we consider the forward orbits of those components of $S^1 \setminus K$ which have to be crossed from the end of β to reach that component. Then there is $N > n$ such that $\bar{s}^{-N} B_0$ intersects none of these components of $S^1 \setminus K$. (Note that, by hypothesis, all components of B_0 lie in strictly preperiodic components of $S^1 \setminus K$.) Take β_k with endpoint close to the same side of L' for all $k < N$. Then, by the choice of N , we can choose β_N having the same intersections with A_{N-1} as β_{N-1} , so that β_N does not intersect the forward orbit of B_N and $\beta_N(1)$ is close to a component of A_N in the minor gap. Then we take the end of β_k close to this component of A_N for all $k \geq N$.

5.7. Moving to a Primitive Lamination.

In Chapter 7 of [R2] we showed how to modify a lamination with eventually periodic minor gap (or leaf) to a primitive lamination. The corresponding lamination maps are then equivalent, and if one has a nondegenerate Levy cycle, they both do. Thus, by the Nonrational Lamination Map Theorem, if one has a piecewise-geodesic periodic holed-sphere, they both do. In fact, it is relatively easy to obtain the holed sphere for one directly from the other if one remembers the definition of primitive, but we omit the details.

5.8. Admissible Points.

The definition of admissible given in 1.17 of [R2] was slightly incorrect. There are several possible rather similar definitions. We give one here which avoids certain complications in the subsequent proof. The definition depends on a parameter lamination $\mathcal{L} = \mathcal{L}_r$ on $\tilde{U}$, in which all leaves are either minor leaves or sides of minor gaps of invariant laminations containing all but countably many leaves of $\bar{L}_r$.

Types II, III and IV. Let f be a degree two branched covering with critical points c_1 and c_2 , with c_1 periodic. Recall from [R2] 1.2 that f is type II if c_2 is in the same periodic orbit as c_1 , type III if c_2 is in the full orbit of c_1 but not periodic, and type IV if c_2 is in a distinct periodic orbit. This includes all critically finite degree two branched coverings such that every critical point has a periodic critical point in its forward orbit.

A leaf $\tilde{\mu}$ in $\mathcal{L}$ is *admissible* if it has a neighbourhood V such that, whenever L is a primitive invariant lamination with minor gap intersecting V , then ρ_L does not have a nondegenerate Levy cycle. Note that if ρ_L is type II, III or IV and has no nondegenerate Levy cycle, then ρ_L is equivalent to a rational map.

A point in $\text{Interior}(\tilde{U})$ is *admissible* if it is either in an admissible leaf of $\mathcal{L}$ or in a gap with at least one admissible side. A point in $\partial\tilde{U}$ is *admissible* if it has a neighbourhood whose intersection with $\text{Interior}(\tilde{U})$ consists of admissible points. The set of admissible points is open by definition, and is denoted by $\tilde{U}_{ad}$.

5.9.

We are now ready to prove the Admissible Boundary Theorem, which we restate. See also the introduction of [R2]. Again, a very slight correction has been made. Recall that if m is the period of r under $x \mapsto 2x \bmod 1$, then m is also the period of 0 under $\bar{s} = \bar{s}_r$, and under lamination maps of all invariant laminations with minor leaves and minor gap sides in $\mathcal{L}_r$.

Admissible Boundary Theorem. *Let $\tilde{\nu}$ be a leaf in $\partial\tilde{U}_{ad}$. Then either $\tilde{\nu}$ is a side of the minor gap $G(L)$ of a primitive invariant lamination L , or there is a sequence $\{L_n\}$ of primitive invariant laminations with minor gaps $G(L_n)$ such that $\tilde{\nu} = \lim_{n \rightarrow \infty} G(L_n)$. Also, the following hold. Let $\rho = \rho_L$ or ρ_{L_n} be the lamination map of L or L_n . Then ρ is type IV, not equivalent to a rational map, and ∞ is periodic of period $\leq m$ under ρ . There is a piecewise-geodesic fixed holed sphere $A_0 \cup B_0$ for L (or L_n) such that A_0 intersects sides of the minor gap of L (or L_n).*

Proof: There are minor gaps of primitive invariant laminations intersecting any neighbourhood of $\tilde{\nu}$ for which some of the corresponding lamination maps have nondegenerate Levy cycles, and some do not. So at least L or $\{L_n\}$ exist with minor gaps $G(L)$, $G(L_n)$ such that either $\tilde{\nu}$ is a side of $G(L)$ or $\tilde{\nu} = \lim_{n \rightarrow \infty} G(L_n)$, and ρ has a nondegenerate Levy cycle, where $\rho = \rho_L$ or ρ_{L_n} . In what follows, the point to remember is that two points on disjoint geodesics are close if the two geodesics have the same S^1 -crossings for a considerable distance in both directions.

Write $\tilde{\mu}$ for the minor leaf of L_n , or $\tilde{\mu} = \tilde{\nu}$ if $\tilde{\nu}$ is a side of $G(L)$, so that $\tilde{\nu}$ is approximated by leaves $\tilde{\mu}$. Write L rather than L_n for the moment. The Nonrational Lamination Map Theorem implies that there is a piecewise-geodesic periodic holed-sphere $A_0 \cup B_0$ for L . Let $A_k \cup B_k$ denote the subset of $\rho^{-k}(A_0 \cup B_0)$ which is homotopic to $A_0 \cup B_0$ in $\bar{\mathbf{C}} \setminus X(\rho)$ and such that each component of A_k contains a component of A_0 . (See 4.2.) The First Invariant-implies-Parameter Theorem can be interpreted as follows. Here, ε , δ depend only on a neighbourhood V of a compact subset of $\tilde{\nu}$ which $G(L)$ intersects. Write $L(A_k, \beta)$ for the lamination L constructed from (A_k, β) as in the First Theorem.

Given $\varepsilon > 0$, there is $\delta > 0$ such that, either $\tilde{\mu} \cap V$ lies in an ε -neighbourhood of $G(L(A_k, \beta))$, or no A_k passes within δ of $V \cap \tilde{\mu}$.

In fact, given $\varepsilon > 0$, the first possibility must hold for some $\tilde{\mu}$. Otherwise, the Second Invariant-implies-Parameter Theorem gives the following.

There is $\eta > 0$, which depends on δ , and hence on ε , such that every primitive lamination L' with minor gap passing within η of $\tilde{\mu} \cap V$ satisfies one of the following.

- a) A_k (for any k) is contained in leaves or finite sided gaps of L' .*
- b) $L' \subset L''$ with $A_k \subset L''$ such that L'' and L' have the same minor leaf.*
- Xc) L' has a nonsimply-connected minor gap.*

By Lemma 5.6 and 5.7, we can choose L'' so that $A_k \cup B_k$ is a piecewise-geodesic periodic holed sphere for L'' , and then modify to a primitive lamination. (The hypothesis of 5.6 is satisfied, because $A_0 \cup B_0$ is a piecewise-geodesic periodic holed-sphere for L .) If $\tilde{\mu}$ is sufficiently close to $\tilde{\nu}$, this implies $\tilde{\nu}$ is in the interior of $\tilde{U} \setminus \tilde{U}_{ad}$, giving a contradiction.

Now using laminations $L(A_k, \beta)$ instead of the original L_n (or L), the corresponding lamination maps are all type IV, not equivalent to rational maps. If the other property (concerning the piecewise-geodesic *fixed* holed-sphere) is not satisfied, then the period of ∞ , for all the maps in the sequence, is less than m , by the Nonrational Lamination Map Theorem. We can then repeat the entire process. After finitely many repeats, we obtain a sequence with the required properties.

5.10. Preliminaries to the Admissibility Proposition.

Every leaf intersecting the admissible set is contained in it. If a lamination map ρ_L of a lamination L is critically finite and equivalent to a rational map, one would expect all sides of the minor gap to be in the admissible set. Actually, if ρ_L is equivalent to a polynomial this cannot quite be true - because then some tunings of ρ_L (with minor gaps contained in the minor gap of L) will not be equivalent to rational maps. But if ρ_L is equivalent to a polynomial and hence to $\mathfrak{F}_r$ for some r , then ρ_L is associated to the gap of $\mathcal{L}_r$ containing 0, which is admissible. Perhaps the easiest degree two rational maps to study are matings and captures, originally defined by Douady and Hubbard [D] and Wittner [W]. They are also described in 1.19 of [R2], where a subset $\tilde{U}_{mc}$ of $\tilde{U}$ is described in which all matings and captures are represented, and a description is given (without proof) of $\tilde{U}_{ad} \cap \tilde{U}_{mc}$. This set is connected and contains up to equivalence all matings and captures which are equivalent to rational maps. For matings, the description is the result of Tan Lei's thesis [TL]. In both cases, it follows from the more general Admissible Boundary Theorem, or even the Nonrational Lamination Map Theorem. A simplifying factor is that, if $\tilde{\mu}$ and $\tilde{\mu}'$ are two minor leaves in $\tilde{U}_{mc}$ of laminations L and L' , and $\tilde{\mu}$ separates $\tilde{\mu}'$ from 0, then all periodic leaves in L are in L' . So now we state the Admissibility Proposition, in a slightly different, but equivalent form to that given in 1.17 of

[R2], given that the definition of admissibility has been altered in 5.8 from that of 1.17 of [R2].

The Admissibility Proposition. *Let f be a critically finite degree two rational map which is not equivalent to a polynomial, mating or capture. Then whenever L is an invariant lamination with minor leaf or gap in $\tilde{U}$ and the lamination map ρ_L is equivalent to f , the minor leaf or gap lies in the admissibility set $\tilde{U}_{ad}$.*

5.11. The Inadmissible Type II Case.

Lemma.

Let L be an invariant lamination with type II lamination map ρ_L which is equivalent to a rational map f . Let L' be the lamination $L \cup \bar{L}_r$ with type IV lamination map ρ' which is not equivalent to a rational map. Then ρ' is equivalent to a mating and ρ_L is equivalent to a capture.

Proof: Let G_∞ be the gap of L containing ∞ , and let $\tilde{G}$ be a lift to the universal cover of $\bar{\mathcal{C}} \setminus K$, with projection $\pi : \tilde{G} \rightarrow \bar{G}_\infty$. Let $\tilde{\rho}$ be a lift of ρ_L^m to G_∞ fixing $\tilde{G}$, and extending to $\tilde{G}$, if m is the period of G_∞ under ρ_L . Let $\Phi : \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$ be the semiconjugacy satisfying $\Phi \circ \rho_L = f \circ \Phi$. Let $\varphi_1 : \{z : |z| \leq 1\} \rightarrow \overline{U_2(f)}$ and $\varphi_2 : \tilde{G} \rightarrow \{z : |z| \leq 1\}$ satisfy

$$\begin{aligned}\varphi_1 \circ \varphi_2 &= \Phi \circ \pi, \\ \varphi_1(z^4) &= f \circ \varphi_1(z), \\ \varphi_2 \circ \tilde{\rho}(z) &= (\varphi_2(z))^4\end{aligned}$$

for all z . Let ω_1, ω_2 be the cube roots of unity which are the images under φ_2 of the endpoints of the lift to $\tilde{G}$ of ℓ , where ℓ is the unique leaf of $\bar{L}_r \setminus L$ of oriented period m in G_∞ . Let α be an arc in $\overline{U_2(f)}$ which is the image under φ_1 of an arc in $\{z : |z| \leq 1\}$ joining ω_1 and ω_2 . We can choose α , and can perturb f to a critically finite branched covering f' , so that $f' = f$ outside a neighbourhood of the critical orbit of f , f' preserves the set $\cup_{i=0}^{m-1} f^i \alpha$, $f^i \alpha \subset f^i U_2(f) = f^i U_2(f)$ separates the two points of $X(f')$ in $f^i U_2(f)$, and $f' \simeq \rho'$. So f' admits a minimal Levy cycle Γ , that is, one with only one complementary component which is not a disc. (See 1.6 of [R2].) Let P be the closure of the nondisc component of $\bar{\mathcal{C}} \setminus (\cup \Gamma)$. Let $F_0 : P \times [0, 1] \rightarrow \bar{\mathcal{C}} \setminus X(f')$ be a homotopy between P and a component of $f'^{-1}(P)$ with $f' \circ F_0(x, 1) = F_0(x, 0)$. (This exists for a minimal Levy cycle.) We can assume $F_0(P \times [0, 1]) \cap (\cup_{i \geq 0} f^i(U_2(f))) \subset \cup_{i \geq 0} f^i \alpha$. Defining F_n inductively by $f' \circ F_{n+1} = F_n$, $F_{n+1}(x, 0) = F_n(x, 1)$, and taking limits, we can assume $\cup \Gamma \subset \cup_{i \geq 0} \overline{f^i \alpha}$, if we drop the assumption that the loops of

Γ are embedded and disjoint. (See Section 8 of [R1] for a similar construction.) This set must contain at least one point fixed by f' by the Lefschetz Theorem. So one endpoint of α must be fixed by f and by f' . The endpoints of α must be distinct if $m \geq 3$, for topological reasons, but also if $m = 2$, since then f is $z \mapsto z^{-2}$, up to Möbius conjugation. Then the second endpoint must also be fixed. Then we can find a circle Γ' invariant up to homotopy under f' , and separating the critical orbits, which is equivalent to f' being a mating. See Diagram 5.

Diagram 5

Then

$$f \simeq \sigma_{\bar{\eta}} \circ \sigma_{\gamma} \circ f',$$

where γ, η are loops crossing Γ' once each, (where $\bar{\eta}$, as usual, is η reversed), and $f'(\eta)$ is homotopic to γ in $\bar{\mathcal{C}} \setminus X(f')$. Let f_0 be the polynomial obtained by modifying f' on the component of $\bar{\mathcal{C}} \setminus \Gamma'$ containing $c_2(f')$, so that $c_2(f_0)$ is fixed. Up to homotopy in $\bar{\mathcal{C}} \setminus X(f_0)$, f_0^{-1} preserves Γ' and the components of $D \setminus (\cup (f')^i \alpha)$, where D is the component of $\bar{\mathcal{C}} \setminus \Gamma$ containing c_1 . Hence, f_0 is a polynomial (corresponding to a minimal minor leaf in *QML*: see 1.10 of [R2]). Then, modifying γ and η also, there are paths β and ζ in $\bar{\mathcal{C}} \setminus X(f_0)$ with $f_0(\zeta)$ homotopic in $\bar{\mathcal{C}} \setminus X(f_0)$ and

$$f \simeq \sigma_{\bar{\zeta}} \circ \sigma_{\beta} \circ f_0,$$

as required.

5.12. Definition of W_1 , W_2 .

Let f be a critically finite rational map with numbered critical points c_1, c_2 . Let c_1 be periodic, and suppose f is not type II. Then we define sets W_1, W_2 as follows. If f is not type III or IV (so that, in particular, c_2 is not periodic) then $W_1 = W_2$ is simply $\{f^i(c_2) : i > 0\}$. If f is type III, then $W_1 = W_1(x) = W_2$ is $\{f^i(x) : i \geq 0\}$ for an eventually periodic point $x \in \partial f(U_2)$. If f is type IV, then W_2 is the closure of the (finite) union of components of $\bar{\mathbf{C}} \setminus J(f)$ which intersect the forward orbit of c_2 , and W_1 is the smallest set with the following properties.

1. $W_2 \subset W_1$
2. W_1 contains $\bar{U}$, whenever U is a component of $\bar{\mathbf{C}} \setminus J(f)$ in the full orbit of $U_2(f)$ with $\bar{U} \cap W_1 \neq \emptyset$.

Then W_1 has finitely many components. By 5.4 of [R2], if U and V are in the same period p orbit of components of $\bar{\mathbf{C}} \setminus J(f)$, with $U \neq V$, then $\bar{U} \cap \bar{V}$ consists of at most one point, which is fixed by f^p .

5.13. Lemma.

Let f be type IV. All components of $\bar{W}_1$ are simply connected, unless f is equivalent to a polynomial or mating

Proof: Suppose that components of $\bar{W}_1$ are not simply connected and that f is not a polynomial, so that $c_2(f)$ has period $p > 1$. By 5.4 of [R2], $\bar{U}$ is simply connected for any component U of $\bar{\mathbf{C}} \setminus J(f)$. So the components of W_1 are larger than the closures of components of $\bar{\mathbf{C}} \setminus J(f)$. There is $x \in \cap_{j \geq 0} f^{jt} \bar{U}_2(f)$ with x of period $t < p$. Let γ be an embedded circle enclosing x and lying in the union of the $f^{jt} U_2(f)$ and a neighbourhood of x , and enclosing x and $f^{jt}(c_2)$ ($j \geq 0$). Then γ is isotopic to a component γ_1 of $f^{-t}(\gamma)$ in $\bar{\mathbf{C}} \setminus J(f)$, with $f^t \mid \gamma_1$ degree 2. Then f must be a tuning $g \vdash_{c_2} h$ of a type IV rational map g (see [R2] 1.20), where h is a polynomial for which the boundaries of the attractive basins have a common fixed point. It is possible to define a branched covering f_1 which is equivalent to f and equal to g on $J(g)$. Moreover, the Semiconjugacy Proposition of 4.1 of [R2] implies that there is $\varphi : \bar{\mathbf{C}} \rightarrow \bar{\mathbf{C}}$ with $\varphi \circ f_1 = f \circ \varphi$, and φ is a limit of homeomorphisms. Since $f_1 \mid J(g)$ is expanding, each set $\varphi^{-1}x$, which is connected and simply-connected, has finite intersection with $J(g)$. (See 4.6 of [R2].) Then $\bar{W}_1 = \varphi(\cup_{i \geq 0} g^i \bar{U}_2(g))$. So components of $\cup_{i \geq 0} g^i \bar{U}_2(g)$ are not simply connected. Then by 5.4 of [R2], g is a polynomial. Then f is a mating, since the tuning of a polynomial is a mating.

5.14. Lemma.

There is an arbitrarily small neighbourhood W of W_1 and such that, for all n , each component of $f^{-n}W$ is either contained in, or disjoint from, W .

Proof: We start with a neighbourhood W' of W_1 such that W' contains the components of $f^{-p}W'$ intersecting W_1 (if p is the least integer > 0 with $f^p W_1 = W_1$). The idea is then to reduce W' to obtain W , without violating the neighbourhood condition. We see that there is $\delta > 0$ with the following property.

a) If W'' is a component of $f^{-n}W'$ for some n , W'' does not contain W_1 and has diameter ε , then $\text{dist}(W_1, W'') > \delta\varepsilon$

Of course, this property remains true if we make W' even smaller. Given any $\eta > 0$, we can choose W' so close to W_1 that the following is true.

b) If W'' is as in a), and $W'' \cap W' \neq \emptyset$, then $\varepsilon < \eta$.

There is $M > 0$ such that the following is true (and remains true if W' is made smaller).

c) For all z, w in the domain of a local inverse S of f^n defined on a subset of $W' \setminus W_1$,

$$\frac{1}{M} \leq \frac{|S'(z)|}{|S'(w)|} \leq M.$$

It follows that if W' is close enough to W_1 , then given $\nu > 0$,

d) if W', W'' are as in a) and $W' \cap W'' \neq \emptyset$ and W'_n, W''_n are preimages under f^n with $W'_n \cap W''_n \neq \emptyset$, then W''_n is contained in the $\text{diameter}(W'_n) \times \nu$ -neighbourhood of W'_n and has diameter $< \text{diameter}(W'_n) \times \nu$.

Now we inductively define $T_0 = W'$, and T_{n+1} is $T_n \setminus R_n$, where R_n is the union of those components of $f^{-k}T_n$ (for all k) which intersect, but are not contained in, T_n . We take $W = \cap_n T_n$. Then W has the required properties - provided W' is close enough to W_1 - because every component of R_{n+1} intersects a component of R_n . Hence we see inductively that every component of R_n is distance $\leq$

$$\sum_{i=1}^n \nu^i \times \text{diameter}(C)$$

from a component C of R_0 which is itself distance $> \delta \times \text{diameter}(C)$ from W_2 , by a). So we only need to take $\frac{\nu}{1-\nu} < \delta$. By its definition T_n contains the component of $f^{-p}(T_n)$ containing W_1 , so W contains the component of $f^{-p}W$ containing W_1 .

5.15. Proof of the Admissibility Proposition.

Let f be a critically finite rational map which is not equivalent to a polynomial, mating or capture. Let L be an invariant lamination with minor leaf or gap in $\tilde{U}$, such that the lamination map ρ is equivalent to f . By 5.11, if f is type II, the lamination map for the lamination $L \cup \bar{L}_r$ is equivalent to a rational map of type IV. So now we can assume that f is not type II. Then the

Semiconjugacy Proposition 4.1 of [R2] gives a semiconjugacy $\Phi : \bar{\mathcal{C}} \rightarrow \bar{\mathcal{C}}$ such that $\Phi \circ \rho = f \circ \Phi$, Φ is a limit of homeomorphisms and collapses each leaf and finite sided gap of L to a point. If f is type III, let x be the image under Φ of the minor leaf of L . Then let W_1 be as in 5.12, with $W_1 = W_1(x)$ if f is type III. By 5.13, $\bar{W}_1$ is simply-connected. Let W be as in 5.14, and we can choose W close enough to W_1 so that $\bar{W}$ is contractible in $\bar{\mathcal{C}} \setminus \{f^i(c_1) : i \geq 0\}$. Put

$$V = \pi^{-1}((\bar{\mathcal{C}} \setminus (K \cup (\cup \bar{L}_r))) \cap \Phi^{-1}W) \subset \tilde{U},$$

where π is the universal covering map. Then $\pi(\bar{V})$ is contractible in $\bar{\mathcal{C}} \setminus \{\bar{s}^i(0) : i \geq 0\}$ and is a neighbourhood of the minor leaf of L , and of the minor gap if this exists, such that $\pi(V)$ contains any component of $\rho^{-n}(V)$ it intersects, for any $n \geq 0$. We claim that the proof of the Admissibility Proposition is completed by the following lemma, using the notation established here.

Lemma. *Let an integer $m > 0$ be given. Let A denote the set of leaves of L of period $\leq m$. Let $V' \subset \mathcal{V}$ be a neighbourhood of the minor leaf or minor gap. Let L' be any primitive invariant lamination with minor gap intersecting V' . Let A' be the set of leaves of L' of period $\leq m$. Then if V' is sufficiently small, every leaf of A' is in A , or a finite-sided gap of L bounded by leaves of A , or in $\pi(V)$.*

We see that the lemma implies the proposition as follows. Let m be the period of r under $x \mapsto 2x \bmod 1$, so that m is the period of 0 under $\bar{s}_r = \bar{s}$, ρ , ρ' , and, indeed, under τ , where τ is any lamination map of an invariant lamination with minor leaf in $\mathcal{L}_r$. Suppose the lamination map ρ' of L' is a critically finite branched covering. By the Nonrational Lamination Map Theorem, ρ' does not have a nondegenerate Levy cycle provided $\bar{A}'$ is contractible in $\bar{\mathcal{C}} \setminus X(\bar{s})$ (to finitely many points). But this is true, because $\overline{A \cup \pi(V)}$ is contractible in $\bar{\mathcal{C}} \setminus X(\bar{s})$, since A maps to points under Φ and $\overline{\pi(V)}$ maps to $\bar{W}$.

5.16. Proof of Lemma 5.15.

Let C be an orbit in A' . We have $C = \cup_{n \geq 0} C_n$, where $C_n \subset C_{n+1} \subset \rho'^{-1}C_n$ and the number of segments in C_0 can be bounded independently of L' . (The bound depends on V' .) By taking V' sufficiently small, given t , we can assume that any geodesic of L of $\geq t$ segments intersecting $\pi(V')$ cannot be isotoped into $\partial\pi(V)$ by good isotopy. We can also take V' small enough that any such C_0 does not lift to lie strictly between the minor leaves of gaps of L , L' , up to good isotopy. Take any k such that C_i does not intersect $\pi(V')$ for $i < k$. By the Second Invariant-implies-Parameter Theorem (with L replacing L'), C_k does not intersect L transversally. So either $C_k \subset \pi(V)$, or $C_k \cap \pi(V) = \phi$, and then C_k lies in leaves or finite-sided gaps of L up to good isotopy. But $\rho^{-1} = \rho'^{-1}$ outside $\pi(V)$, and $\rho^{-k}(\pi(V)) = \rho'^{-k}(\pi(V))$ for all $k \geq 0$. So either

$C \subset \cup_{n \geq 0} \rho'^{-n} C_0$ and $C \subset \pi(V)$ or, for all k , C_k lies in leaves or finite-sided gaps of $\bar{L}$ up to good isotopy, and hence so does C .

References

- [D] Douady, A.: Systèmes dynamiques holomorphes. Séminaire Bourbaki 1983, *Astérisque* 105/6 (1983), 39-63.
- [D-H1] Douady, A. and Hubbard, J.H.: Etudes dynamiques des polynômes complexes, avec la collaboration de P. Lavaurs, Tan Lei, P. Sentenac. Parts I and II, Publications Mathématiques d'Orsay, 1985.
- [D-H2] Douady, A. and Hubbard, J.H.: Itérations des polynômes quadratiques complexes. *C.R.Acad.Sci. Paris, Série I, t.294* (1982), 123-126.
- [D-H3] Douady, A. and Hubbard, J.H.: A proof of Thurston's topological characterization of rational functions. Mittag-Leffler preprint, 1985.
- [F-L-P] Fahti, A., Laudenbach, F., and Poénaru, V.: Travaux de Thurston sur les Surfaces. *Astérisque* 66-67 (1989).
- [Fr] Franks, J.: Anosov Diffeomorphisms. *Proc. Symposia in Pure Math.*, 14 (1968), 61-93.
- [R1] Rees, M.: Components of degree two hyperbolic rational maps. *Invent. Math.*, 100 (1990), 357-382.
- [R2] Rees, M.: A Partial Description of the Parameter Space of Rational Maps of Degree Two. Part 1. To appear in *Acta Math.*
- [S] Shishikura, M.: In preparation.
- [T] Thurston, W.P.: On the Combinatorics of Iterated Rational Maps. Preprint, Princeton University and I.A.S., 1985.
- [TL] Tan Lei: Accouplements des polynômes complexes. Thèse, Université de Paris-Sud, Orsay, 1987.
- [W] Wittner, B.: On the bifurcation loci of rational maps of degree two. Thesis, Cornell University, 1988.